


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1977

[Full Table of Content](#)



77/W/RN/13 Intensive Cereals - Wheat, Barley

Rothamsted Research

Rothamsted Research (1978) *77/W/RN/13 Intensive Cereals - Wheat, Barley ; Yields Of The Field Experiments 1977*, pp 85 - 88 - DOI: <https://doi.org/10.23637/ERADOC-1-29>

77/W/RN/13

INTENSIVE CEREALS

Object: To study the effects of intensive cereal cropping on yield, incidence of soil-borne pathogens and organic matter in the soil - Woburn Stackyard I.

Sponsors: A.E. Johnston, J. McEwen.

The 12th year, winter wheat, barley.

For previous years see 'Details' 1973 and 74-76/W/RN/13.

Design: For each experiment: 2 randomised blocks of 6 plots, split into 4. ALDICARB tested on blocks.

Whole plot dimensions: 8.53 x 20.4.

Treatments:-

One experiment on winter wheat on part of the site of the classical wheat experiment 1877-1954

One experiment on barley on part of the site of the classical barley experiment 1877-1954

Factors tested on both experiments are the same but crop and nitrogen rates differ. All combinations of:-

Blocks

1. ALDICARB Aldicarb worked into the seedbed (kg):

0
10

Whole plots

2. PREVCROP Previous crops:

	1972	1973	1974	1975	1976
C/C/L/P	C	C	C	L	P
C/L/P/C	C	C	L	P	C
L/P/C/C	C	L	P	C	C
P/C/C/C	L	P	C	C	C
C/C/C/L	P	C	C	C	L
C/C/C/C	C	C	C	C	C

Ley = 1 year ley P = Potatoes C = Cereal: wheat or barley. All plots in cereal only from 1977.

Sub plots

3. N Nitrogen fertiliser (kg N):

Wheat	Barley
62	50
126	100
189	150
252	200

77/W/RN/13

Standard applications:

Wheat: Manures: (0:20:20) at 290 kg, combine drilled. Weedkillers: Glyphosate at 1.7 kg in 280 l, methabenzthiazuron at 1.6 kg in 280 l, Ioxynil at 0.53 kg plus mecoprop at 1.6 kg in 220 l. Insecticide: Pirimicarb at 0.14 kg in 270 l.

Barley: Manures: (0:20:20) at 300 kg, combine drilled. Weedkillers: Glyphosate at 1.7 kg in 280 l, Ioxynil at 0.53 kg plus mecoprop at 1.6 kg in 220 l.

Seed:

Wheat: Cappelle, sown at 210 kg.

Barley: Julia, dressed with ethirimol, sown at 160 kg.

Cultivations, etc.:— All plots: Glyphosate applied: 24 Sept, 1976. Rotary cultivated grass leys: 1 Nov. Ploughed: 8 Nov.

Wheat: Aldicarb applied, rotary cultivated, seed sown: 24 Nov, 1976. Methabenzthiazuron applied: 25 Nov. N applied: 13 Apr, 1977. Ioxynil plus mecoprop applied: 15 May. Pirimicarb applied: 11 July. Combine harvested: 7 Sept.

Barley: Spring-tine cultivated: 7 Mar, 1977. Aldicarb applied, rotary cultivated, seed sown: 8 Apr. N applied: 13 Apr. Ioxynil plus mecoprop applied: 19 May. The following treatments were combine harvested on 16 Aug:

ALDICARB 0	ALDICARB 10	ALDICARB 0	ALDICARB 10
PREVCROP C/C/C/L	PREVCROP P/C/C/C	PREVCROP C/C/L/P	PREVCROP C/L/P/C
N 50, 150, 200	N 50, 150	N 100, 200	N 100, 150

Remaining treatments combine harvested: 30 Aug.

77/W/RN/13

WINTER WHEAT

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	C/C/L/P	C/L/P/C	L/P/C/C	P/C/C/C	C/C/C/L	C/C/C/C	MEAN
ALDICARB							
0	3.79	3.58	3.38	2.84	3.44	3.26	3.38
10	3.89	3.65	3.43	2.37	2.70	2.90	3.16
MEAN	3.84	3.62	3.41	2.61	3.07	3.08	3.27

N	63	126	189	252	MEAN
ALDICARB					
0	3.15	3.48	3.56	3.34	3.38
10	2.81	3.55	3.17	3.10	3.16
MEAN	2.98	3.51	3.36	3.22	3.27

N	63	126	189	252	MEAN
PREVCROP					
C/C/L/P	3.55	4.39	3.76	3.66	3.84
C/L/P/C	3.51	4.07	3.60	3.28	3.62
L/P/C/C	2.94	3.50	3.65	3.54	3.41
P/C/C/C	1.98	2.64	2.88	2.93	2.61
C/C/C/L	2.90	3.31	3.00	3.08	3.07
C/C/C/C	3.02	3.18	3.28	2.85	3.08
MEAN	2.98	3.51	3.36	3.22	3.27

		N	63	126	189	252
ALDICARB	PREVCROP					
0	C/C/L/P	3.57	4.10	3.68	3.82	
	C/L/P/C	3.53	3.86	3.51	3.42	
	L/P/C/C	2.79	3.60	3.53	3.59	
	P/C/C/C	2.39	2.72	3.18	3.07	
	C/C/C/L	3.15	3.57	3.80	3.26	
	C/C/C/C	3.50	3.02	3.64	2.90	
10	C/C/L/P	3.53	4.69	3.85	3.49	
	C/L/P/C	3.49	4.28	3.70	3.14	
	L/P/C/C	3.09	3.40	3.76	3.49	
	P/C/C/C	1.57	2.56	2.58	2.79	
	C/C/C/L	2.66	3.05	2.21	2.90	
	C/C/C/C	2.55	3.34	2.92	2.81	

GRAIN MEAN DM% 82.6

SUB PLOT AREA HARVESTED 0.00277

77/W/RN/13

BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	C/C/L/P	C/L/P/C	L/P/C/C	P/C/C/C	C/C/C/L	C/C/C/C	MEAN
ALDICARB							
0	4.26	4.36	4.49	4.22	4.53	4.06	4.32
10	4.29	4.74	4.48	4.02	4.31	3.82	4.28
MEAN	4.28	4.55	4.48	4.12	4.42	3.94	4.30

N	50	100	150	200	MEAN
ALDICARB					
0	3.35	4.60	4.67	4.67	4.32
10	2.95	4.65	4.98	4.53	4.28
MEAN	3.15	4.63	4.82	4.60	4.30

N	50	100	150	200	MEAN
PREVCROP					
C/C/L/P	3.15	4.62	4.58	4.76	4.28
C/L/P/C	3.88	5.07	4.97	4.29	4.55
L/P/C/C	3.45	4.81	4.97	4.71	4.48
P/C/C/C	2.82	4.33	5.16	4.19	4.12
C/C/C/L	2.83	4.82	4.98	5.06	4.42
C/C/C/C	2.78	4.11	4.28	4.59	3.94
MEAN	3.15	4.63	4.82	4.60	4.30

N	50	100	150	200
ALDICARB PREVCROP				
0 C/C/L/P	3.49	4.58	4.31	4.67
C/L/P/C	3.70	4.70	4.65	4.40
L/P/C/C	3.85	4.65	4.63	4.82
P/C/C/C	3.38	4.46	4.72	4.34
C/C/C/L	2.78	4.94	5.05	5.34
C/C/C/C	2.91	4.27	4.63	4.42
10 C/C/L/P	2.82	4.66	4.84	4.86
C/L/P/C	4.06	5.43	5.29	4.17
L/P/C/C	3.05	4.97	5.31	4.60
P/C/C/C	2.26	4.20	5.60	4.04
C/C/C/L	2.87	4.70	4.90	4.78
C/C/C/C	2.64	3.95	3.93	4.76

GRAIN MEAN DM% 81.7

SUB PLOT AREA HARVESTED 0.00277