

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1978

[Full Table of Content](#)



78/R/CS/24 P K and Take-all - Wheat

Rothamsted Research

Rothamsted Research (1979) *78/R/CS/24 P K and Take-all - Wheat ; Yields Of The Field Experiments 1978*, pp 144 - 146 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/R/CS/24

PK AND TAKE-ALL

Object: To study the effects of different amounts of phosphate and potassium fertiliser on the yields and incidence of take-all (*Gaeumannomyces graminis*) in continuous wheat - West Barnfield II.

Sponsors: G.E.G. Mattingly, D.B. Slope.

The 11th year, continuous winter wheat (after continuous barley 1968-1973).

For previous years see 'Details' 1973 and 74-77/R/CS/24.

Design: 4 randomised blocks of 10 plots, split into 2.

Whole plot dimensions: 5.33 x 20.1.

Treatments: All combinations of:-

Whole plots

1. P Phosphate (kg P) as superphosphate:

0	None
15 A	15 annually
60 A	60 annually
90 S	90 six-yearly, last applied autumn 1973
360 S	360 six-yearly, last applied autumn 1973

2. K Potassium (kg K) annually as muriate of potash:

30
120

Sub plots

3. N Nitrogen fertiliser, applied cumulatively to test applications 1970-1973 (basal application only in 1974-1977) (kg N):

50
100
150
200

Basal applications: Autumn weedkiller: Glyphosate at 1.7 kg in 220 l. Spring weedkillers: Mecoprop, bromoxynil and ioxynil ('Brittox' at 2.8 l in 220 l).

Seed: Cappelle sown at 200 kg.

Cultivations, etc.: Autumn weedkiller applied: 10 Oct, 1977. Ploughed: 25 Oct. Spring-tine cultivated: 26 Oct. P and K applied, power harrowed, seed sown: 18 Nov. Rolled: 19 Apr, 1978. N applied: 24 Apr. Spring weedkiller applied: 11 May. Combine harvested: 30 Aug.

NOTE: The crop was sampled in July for take-all and eyespot assessments.

78/R/CS/24

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	K	30	120	MEAN
P				
O		3.49	3.57	3.53
15 A		4.20	4.65	4.43
60 A		4.74	5.18	4.96
90 S		4.31	4.55	4.43
360 S		4.64	4.83	4.74
MEAN		4.28	4.56	4.42

	N	50	100	150	200	MEAN
P						
O		3.40	2.98	3.85	3.89	3.53
15 A		3.79	4.73	4.68	4.51	4.43
60 A		4.48	4.86	5.20	5.29	4.96
90 S		3.98	4.57	4.78	4.39	4.43
360 S		4.07	4.92	5.25	4.71	4.74
MEAN		3.94	4.41	4.75	4.56	4.42

	N	50	100	150	200	MEAN
K						
30		3.86	4.37	4.43	4.44	4.28
120		4.03	4.46	5.07	4.67	4.56
MEAN		3.94	4.41	4.75	4.56	4.42

	N	50	100	150	200
P					
O	K				
	30	3.38	3.14	3.48	3.96
	120	3.41	2.83	4.22	3.82
15 A	30	3.98	4.48	4.17	4.18
	120	3.60	4.97	5.19	4.84
60 A	30	4.27	4.49	4.91	5.28
	120	4.68	5.23	5.48	5.30
90 S	30	3.75	4.79	4.43	4.27
	120	4.21	4.35	5.14	4.51
360 S	30	3.91	4.94	5.19	4.54
	120	4.24	4.91	5.31	4.88

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	P	K	N	P K
SED	0.133	0.084	0.119	0.188

TABLE	P N	K N	P K N
SED	0.266	0.170	0.395

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP+BLOCK.WP.SP	37	0.377	8.5

78/R/CS/24

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

K	30	120	MEAN
P			
O	2.39	2.65	2.52
15 A	2.71	3.46	3.08
60 A	3.24	4.08	3.66
90 S	2.70	3.23	2.97
360 S	3.22	3.72	3.47
MEAN	2.85	3.43	3.14

N	50	100	150	200	MEAN
P					
O	2.58	1.95	2.92	2.64	2.52
15 A	2.89	3.42	3.14	2.88	3.08
60 A	3.11	3.62	4.02	3.89	3.66
90 S	2.51	3.03	3.28	3.04	2.97
360 S	2.69	3.57	4.01	3.61	3.47
MEAN	2.76	3.12	3.48	3.21	3.14

N	50	100	150	200	MEAN
K					
30	2.60	2.85	3.00	2.96	2.85
120	2.91	3.39	3.95	3.47	3.43
MEAN	2.76	3.12	3.48	3.21	3.14

P	N	50	100	150	200
K					
O	30	2.37	2.00	2.69	2.51
	120	2.79	1.89	3.15	2.77
15 A	30	3.01	2.96	2.41	2.45
	120	2.76	3.88	3.88	3.30
60 A	30	2.66	3.04	3.69	3.57
	120	3.57	4.20	4.36	4.20
90 S	30	2.46	3.08	2.39	2.89
	120	2.56	2.98	4.18	3.20
360 S	30	2.50	3.17	3.84	3.37
	120	2.89	3.97	4.17	3.86

GRAIN MEAN DM% 84.1

STRAW MEAN DM% 91.0

PLOT AREA HARVESTED 0.00270