

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1978

[Full Table of Content](#)



78/R/CS/90 Cultivations for Cereals - Wheat

Rothamsted Research

Rothamsted Research (1979) *78/R/CS/90 Cultivations for Cereals - Wheat ; Yields Of The Field Experiments 1978*, pp 164 - 165 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/R/CS/90

CULTIVATIONS FOR CEREALS

Object: To study the effects of different tillage systems for continuous wheat - Meadow.

Sponsors: R. Moffitt, R.D. Prew, C.A. Edwards.

The seventh year, winter wheat.

For previous years see 72-77/R/CS/90.

Design: 3 randomised blocks of 4 plots.

Whole plot dimensions: 13.7 x 33.8.

Treatments

TILLAGE Tillage systems:-

- 1 Three passages of the tractor (three-pass system): Ploughed 20 cm deep (8 inches): disc harrowed: drilled.
- 2 Three-pass system: Chisel-ploughed once 15 cm deep (6 inches): disc harrowed: drilled.
- 8 Two-pass system: Sprayed with paraquat (0.56 kg ion in 220 l on 3 Oct, 1977): direct drilled.
- 9 Three-pass system: Rotary digger (N.I.A.E.) cultivated: disc harrowed: drilled.

NOTES: (1) Rotary digger (N.I.A.E.) - depth of working: rotor 10 cm, tines 20 cm.
(2) Treatments 3-7 and 10 were not included in the experiment in 1978.

TILLAGE 1 plots ploughed: 31 Oct, 1977. TILLAGE 2 plots chisel ploughed: 10 Oct.
TILLAGE 9 plots rotary dug: 3 Oct. TILLAGE 1, 2 and 9 disc harrowed, sown, harrowed: 7 Nov. TILLAGE 8 direct drilled and disc harrowed in: 7 Nov.

Basal applications: Manures: (10:24:24) at 250 kg, combine drilled. 'Nitra-Shell 34' at 360 kg. Weedkillers: Paraquat at 0.84 kg ion in 220 l, methabenzthiazuron at 3.1 kg in 220 l, and difenzoquat at 0.99 kg with 1 l 'Agral' (a wetting agent) in 220 l.

Seed: Maris Huntsman, sown at 190 kg.

Cultivations, etc.: Chopped stubble: 19 Sept, 1977. Paraquat applied: 30 Oct. Methabenzthiazuron applied: 11 Nov. N applied: 7 Apr, 1978. Difenzoquat applied: 10 May. Combine harvested: 29 Aug.

NOTES: (1) Disease and pest assessments were made during the season.
(2) Measurements were made on TILLAGE 1 and TILLAGE 8 of the following:-
In the soil: total organic matter, organic carbon, organic nitrogen, microbial bio-mass, total soil adenosine triphosphate. In the plants: total nitrogen at flowering.

78/R/CS/90

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TILLAGE	1	2	8	9	MEAN
	6.40	6.02	4.71	6.44	5.89

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TILLAGE
-----	-----
SED	0.410

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	6	0.502	8.5

GRAIN MEAN DM% 82.9

PLOT AREA HARVESTED 0.01031