


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1978

[Full Table of Content](#)



78/R/CS/211 Factors Affecting Eyespot - Wheat

Rothamsted Research

Rothamsted Research (1979) *78/R/CS/211 Factors Affecting Eyespot - Wheat* ; Yields Of The Field Experiments 1978, pp 290 - 291 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/R/CS/211

FACTORS AFFECTING EYESPOT

Object: To study the effects of a range of treatments on the incidence of eyespot (*Pseudocercospora herpotrichoides* previously *Cercospora herpotrichoides*) and on the yield of wheat - Meadow.

Sponsors: R.D. Prew, A. Bainbridge.

The first year, wheat.

Design: 2 randomised blocks of 2 whole plots split into 6.

Whole plot dimensions: 94.0 x 9.14.

Treatments: All combinations of:-

Whole plots

1. STRAW Treatment of straw in autumn 1977:

BURNT	Burnt on site after spreading
CARTED	Baled and carted off

Sub plots

2. DRILLING Cultivations and drilling:

CNVNTIAL	Cultivated and conventionally drilled	) Treatments
DIRECT	Uncultivated, direct drilled	) triplicated

NOTES: (1) An intended test of three sowing dates on sub plots and three seed rates on sub sub plots was not made this year. Sub sub plots were all sown at 150 kg seed but were separately harvested.
(2) DRILLING CNVNTIAL plots: Ploughed: 31 Oct, 1977. Spring-tine cultivated, seed sown and harrowed in: 7 Nov.
(3) DRILLING DIRECT plots: Seed sown 7 Nov. Chain harrowed in: 8 Nov.
(4) Because of very poor establishment, yields were not taken from one sub plot of treatment combination STRAW CARTED, DRILLING DIRECT. An estimated value was used in the analysis.

Basal applications: Manures: (10:24:24) at 250 kg. 'Nitra-Shell 34' at 360 kg.
Weedkillers: Paraquat at 0.56 kg ion in 220 l. Methabenzthiazuron at 1.3 kg in 220 l.

Seed: Kador, sown at 150 kg.

Cultivations, etc.: - Straw treatments applied: 27 Sept, 1977. Paraquat applied: 11 Oct. Seed sown: 7 Nov. Methabenzthiazuron applied: 11 Nov. N applied: 7 Apr, 1978. Combine harvested: 30 Aug.

NOTE: Emergence counts were made and eyespot was assessed at intervals during the season.

78/R/CS/211

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

DRILLING	CNVNTIAL	DIRECT	MEAN
STRAW			
BURNT	5.05	5.03	5.04
CARTED	5.35	5.58	5.46
MEAN	5.20	5.30	5.25

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	DRILLING	STRAW* DRILLING
SED	0.218	0.309

* WITHIN SAME LEVEL OF STRAW ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	17	0.535	10.2

GRAIN MEAN DM% 84.8

SUB PLOT AREA HARVESTED 0.00257