

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1978

[Full Table of Content](#)



78/R/BE/13 Comparison of Fungicides - Beans

Rothamsted Research

Rothamsted Research (1979) *78/R/BE/13 Comparison of Fungicides - Beans* ; Yields Of The Field Experiments 1978, pp 420 - 421 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/R/BE/13

SPRING BEANS

COMPARISON OF FUNGICIDES

Object: To study the effects of a range of fungicides and methods of application on the incidence of diseases and on yield - Fosters O & E IV.

Sponsors: G.A. Salt, J. McEwen, D. Yeoman.

Design: 3 randomised blocks of 12 plots.

Whole plot dimensions: 2.03 x 2.13.

Treatments: All combinations of:-

1. FUNGICIDE	Fungicide:
AL TRI	Aluminium tris (ethylphosphonate). 'Aliette'
BEN	Benomyl
DL ME	DL-methyl N-(2,6 dimethylphenyl)-N-(2 methoxyacetyl) alaninate. 'Ridomil'
THIABEND	Thiabendazole

2. METHOD	Method of application:
FOLIAR	Foliar spray
SEEDRESS	Seed dressing (using a methyl cellulose sticker)

plus four extra treatments:

EXTRA

NONE	No fungicides
NONE S	No fungicides, seed treated with methyl cellulose sticker
BEN SB	Benomyl worked in to seedbed
DL ME SB	'Ridomil' worked in to seedbed

NOTES: (1) Rates of application were as follows (foliar sprays were applied in 1250 l):

	Seedbed kg	Seed dressing g/kg seed	Foliar spray kg
AL TRI	-	4.0	3.38
BEN	20.0	8.8	0.50
DL ME	1.5	0.4	0.34
THIABEND	-	11.3	0.50

(2) Seed was sown by hand in rows 51 cm apart, seed spaced 5 cm apart in the row.

78/R/BE/13

Basal applications: Weedkillers: Simazine at 0.84 kg in 340 l. Insecticides: Permethrin at 0.15 kg in 340 l. Pirimicarb at 0.14 kg in 340 l.

Seed: Minden.

Cultivations, etc.:— Ploughed: 11 Oct, 1977. Spring-tine cultivated, seedbed treatments applied: 15 Mar, 1978. Rotary cultivated, seed sown: 31 Mar. Simazine applied: 24 Apr. Weedkiller applied: 10 May. Permethrin applied: 17 May. Pirimicarb applied: 5 July. Foliar spray treatments applied: 18 July. Harvested by hand: 11 Oct. Previous crops: Fallow 1976, spring oats 1977.

NOTE: Observations were made on root and foliar diseases during the season.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

METHOD	FOLIAR	SEEDRESS	MEAN
FUNGICIDE			
AL TRI	5.66	5.80	5.73
BEN	5.70	6.00	5.85
DL ME	5.81	5.62	5.71
THIABEND	5.54	4.59	5.07
MEAN	5.68	5.50	5.59

EXTRA	NONE	NONE S	BEN SB	DL ME SB	MEAN
	5.67	5.79	5.39	5.43	5.57

GRAND MEAN 5.58

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	FUNGICIDE	METHOD	FUNGICIDE METHOD & EXTRA
SED	0.441	0.312	0.220	0.441

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	22	0.540	9.7

GRAIN MEAN DM% 80.7

PLOT AREA HARVESTED 0.00015