

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1979

[Full Table of Content](#)



79/R/CS/203 Species Mixtures and Phialophora - Wheat

Rothamsted Research

Rothamsted Research (1980) *79/R/CS/203 Species Mixtures and Phialophora - Wheat ; Yields Of The Field Experiments 1979*, pp 206 - 208 - DOI: <https://doi.org/10.23637/ERADOC-1-45>

79/R/CS/203

SPECIES MIXTURES AND PHIALOPHORA

Object: To study the effects of lucerne, grass and clover, singly and in mixtures, and of soil inoculation on populations of *Phialophora radiculicola* graminicola (Prg) and on take-all (*Gaeumannomyces graminis*) and yield of following wheat crops - Stubbings.

Sponsors: E. Lester, D.B. Slope, R.J. Gutteridge.

The third year, wheat; the first year of yields.

Design: 4 randomised blocks of 9 plots, split into 4.

Whole plot dimensions: 4.27 x 27.1.

Treatments: All combinations of:-

1. Whole plots

CRP INOC	Crops in 1977 and 1978 (all wheat 1979) and inoculation:-	
	1977	1978
C C	White clover	White clover
G G	Ryegrass	Ryegrass
GC GC	Ryegrass/white clover mixture	Ryegrass/white clover
LU LU	Lucerne	Lucerne
LU LU I	Lucerne	Lucerne + Prg inoculum to 1979 wheat
GLU GLU	Ryegrass/lucerne in alternate rows	Ryegrass/lucerne
W G	Spring wheat	Ryegrass, sown into spring wheat stubble
WG G	Spring wheat undersown with ryegrass	Ryegrass
WGI G	Spring wheat, inoculated Prg, undersown with ryegrass	Ryegrass

2. Sub plots

N	Nitrogen fertiliser (kg N) in 1979 only:
0	
50	
100	
150	

NOTES: (1) The inoculum used for the I treatments was an agar culture of Prg mixed with sand. It was broadcast and power harrowed into the soil before sowing.

(2) All lucerne was inoculated with 'Nodulaid'.

79/R/CS/203

Standard applications: 1977 & 1978:

Nitrogen (kg N):	1977	1978
C C and LU LU	0	0
G G	75 per cut	75 per cut
GC GC	30 per cut	30 per cut
GLU GLU	0	30 per cut
W G and WG G	75 to seedbed	75 per cut

NOTE: The first N per cut was applied to herbage crops in the seedbed in the year of sowing, in spring thereafter. Subsequent dressings were applied after each cut except the last.

PK: (0:14:28) at 690 kg to all plots for 1977. None in 1978. Weedkillers: Glyphosate at 1.8 kg in 220 l to all plots for 1977. MCPB (as 'Tropotox' at 7.0 kg in 340 l) to C, GC, and G treatments for 1977. 2,4-DB at 2.5 kg in 340 l to LU and GLU treatments for 1977. None in 1978.

Varieties and seed rates (kg) 1977:

	Grass RvP	Clover Blanca	Lucerne Vertus	S. wheat Sappo
C	-	11	-	-
G	22	-	-	-
GC	22	2.2	-	-
LU	-	-	13	-
GLU	11	-	13	-
W	-	-	-	190
WG	22	-	-	190

NOTE: GC: Grass and clover seeds were mixed. GLU: Grass and lucerne seeds were sown in alternate rows.

Varieties and seed rate 1978: Only grass was sown in 1978; W G: RvP sown at 22 kg.

Cultivations, etc. for 1977:- Glyphosate applied: 20 Sept, 1976. Ploughed: 29 Oct. Spring-tine cultivated: 5 Apr, 1977. PK applied: 6 Apr. Spring-tine cultivated, power harrowed for W and WG only: 7 Apr. Power harrowed for G, rolled G, W and WG: 8 Apr. Seed sown for G, W and WG: 9 Apr. First standard N applied: 19 May. LU, C, GC and GLU power harrowed, G and GC sown: 27 May. LU and GLU sown: 1 June. G and GC cut: 21 July. N applied to G only: 27 July. 'Tropotox' applied: 28 July. 2,4-DB applied: 4 Aug. All wheat cut green: 5 Aug. All plots cut: 8 Nov. Previous cropping: Potatoes 1975, wheat 1976.

Cultivations, etc. for 1978:- W G heavy spring-tine cultivated: 16 Sept, 1977. W G disc harrowed twice: 28 Sept. Seed sown for W G treatments: 3 Oct. N applied: 20 Mar, 1978. Grass cut, N applied (as 'Nitra-Shell 34'): 9 June. Grass cut: 24 July.

Basal applications for 1979: Manures: (0:20:20) at 250 kg, combine drilled. Weedkillers: Paraquat at 0.84 kg ion in 220 l. Mecoprop at 2.5 kg and isoproturon at 2.2 kg in 220 l.

79/R/CS/203

Seed for 1979: Flanders, sown at 200 kg.

Cultivations for 1979: Paraquat applied: 3 Aug, 1978. Heavy spring-tine cultivated: 10 Aug (twice), 25 Aug, 8 Sept, 15 Sept. Power harrowed: 10 Oct. Seed sown: 11 Oct. N applied to half of site: 12 Apr, 1979. N applied to remaining plots: 17 Apr. Isoproturon and mecoprop applied: 9 May. Combine harvested: 30 Aug.

NOTE: Estimates of take-all and Phialophora were made on crop in April and early July. Bio-assays of soils for take-all and Phialophora were made after harvest, before ploughing.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	N	0	50	100	150	MEAN
CRP INOC						
C C		4.46	6.46	7.88	8.28	6.77
G G		3.99	5.49	6.50	6.62	5.65
GC GC		3.64	5.82	6.76	6.86	5.77
LU LU		4.33	6.25	7.43	7.49	6.38
LU LU I		4.13	6.40	7.72	7.78	6.51
GLU GLU		3.47	5.69	6.56	7.70	5.86
W G		3.63	5.41	6.12	6.98	5.54
WG G		3.23	5.29	5.75	6.46	5.18
WGI G		3.55	5.61	6.31	6.61	5.52
MEAN		3.83	5.83	6.78	7.20	5.91

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CRP INOC	N	CRP INOC N
SED	0.264	0.144	0.458
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:			
CRP INOC			0.432

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	24	0.374	6.3
BLOCK.WP.SP	81	0.611	10.3

GRAIN MEAN DM% 86.1

SUB PLOT AREA HARVESTED 0.00127