

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1979

[Full Table of Content](#)



79/R/CS/216 and 79/W/CS/216 Effects of Subsoiling and Deep P K - Beans

Rothamsted Research

Rothamsted Research (1980) *79/R/CS/216 and 79/W/CS/216 Effects of Subsoiling and Deep P K - Beans* ; Yields Of The Field Experiments 1979, pp 224 - 225 - DOI:
<https://doi.org/10.23637/ERADOC-1-45>

79/R/CS/216 and 79/W/CS/216

EFFECTS OF SUBSOILING & DEEP PK

Object: To study the effects of subsoiling and of incorporating a large dressing of PK in the subsoil on yields and nutrient uptakes of a sequence of crops - Rothamsted (R) Delharding and Woburn (W) Road Piece.

Sponsors: J. McEwen, A.E. Johnston (R), M.K.V. Carr, R.J. Godwin (National College of Agricultural Engineering), P.T. Gooderham, I.B. Warboys, J.M. Wilkes (Wye College).

The second year, beans.

For previous year see 78/R&W/CS/216.

Design: 3 randomised blocks of 6 plots.

Whole plot dimensions: 4.27 x 13.7.

Treatments (applied autumn 1977 only, before spring barley 1978):

TREATMNT	Machines and incorporation of P and K into the subsoil:
NONE	Not subsoiled, no P or K
FARM O	Farm standard, unwinged, subsoiler, no P or K
NCAE O	N.C.A.E. winged subsoiler, no P or K
NCAE PK	N.C.A.E. winged subsoiler, P and K applied
WYE O	Wye double digger, no P or K
WYE PK	Wye double digger, P and K applied

- NOTES: (1) The rates of P and K were 1930 kg P_2O_5 , as triple superphosphate and 460 kg K_2O as muriate of potash.
- (2) The Farm standard, unwinged, subsoiler was set to work at a depth of 38 cm at intervals of 50 cm Delharding (R) and at a depth of 50 cm at intervals of 70 cm Road Piece (W).
- (3) The N.C.A.E. winged subsoiler was set to work at a depth of 40 cm at intervals of 60 cm on plots not given P and K and at alternate depths of 30 cm and 40 cm spaced 30 cm apart on plots given P and K. Fertiliser was applied behind the subsoiling points.
- (4) The Wye double digger turns a furrow with a conventional plough to a depth of 23 cm, and at the same time rotary cultivates the bottom of the furrow to a further depth of 15 cm. When applying P and K this was distributed ahead of the rotary cultivator.

Basal applications:

Delharding (R): Weedkillers: Glyphosate at 1.5 kg in 220 l. Trietazine with simazine ('Rental SC' at 2.8 kg in 220 l). Insecticide: Pirimicarb at 0.14 kg in 220 l.

Road Piece (W): Weedkillers: Diquat at 0.79 kg in 280 l. Trietazine with simazine ('Rental SC' at 2.4 kg in 280 l). Insecticide: Pirimicarb at 0.14 kg in 250 l.

Seed: Minden, sown at 220 kg on both sites.

Cultivations, etc.:-

Delharding (R): Glyphosate applied: 24 Oct, 1978. Ploughed: 16 Nov. Heavy spring-tine cultivated: 20 Apr, 1979. Rotary harrowed: 21 Apr. Seed sown: 23 Apr. Trietazine with simazine applied: 13 May. Insecticide applied: 22 June. Combine harvested: 21 Sept.

79/R/CS/216 and 79/W/CS/216

Road Piece (W): Diquat applied: 8 Sept, 1978. Heavy spring-tine cultivated: 13 Sept. Ploughed: 16 Nov. Spring-tine cultivated with crumbler attached: 16 Apr, 1979. Seed sown: 19 Apr. Trietazine with simazine applied: 1 May. Insecticide applied: 22 June. Combine harvested: 10 Sept.

NOTES: (1) On Road Piece (W) water use was measured during the season using a neutron probe.
(2) On both sites samples were taken at intervals during the season to measure above-ground dry matter and nutrient uptake. Grain samples were analysed for N, P, K, Ca, Na and Mg.

79/R/CS/216 DELHARDING (R)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	NONE	FARM O	NCAE O	NCAE PK	WYE O	WYE PK	MEAN
	3.61	3.61	3.66	3.49	3.76	4.10	3.71

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT
SED	0.201

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	10	0.247	6.7

GRAIN MEAN DM% 80.3 PLOT AREA HARVESTED 0.00300

79/W/CS/216 ROAD PIECE (W)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	NONE	FARM O	NCAE O	NCAE PK	WYE O	WYE PK	MEAN
	1.41	1.22	1.33	1.29	1.52	1.89	1.44

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT
SED	0.375

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	10	0.460	31.9

GRAIN MEAN DM% 87.3 PLOT AREA HARVESTED 0.00260