

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1979

[Full Table of Content](#)



79/R/WW/5 Fungicides - W. Wheat

Rothamsted Research

Rothamsted Research (1980) *79/R/WW/5 Fungicides - W. Wheat ; Yields Of The Field Experiments 1979*, pp 273 - 274 - DOI: <https://doi.org/10.23637/ERADOC-1-45>

79/R/WW/5

WINTER WHEAT

FUNGICIDES

Object: To study the effects of a range of fungicides and two methods of application on the incidence of diseases and on the yield of winter wheat - Webbs.

Sponsors: J.F. Jenkyn, R.D. Prew.

Design: 3 randomised blocks of 14 plots.

Whole plot dimensions: 2.13 x 13.4.

Treatments:

FUNGICIDE	Fungicides and methods of application:
NONE	None
BAS389 S	'BAS 389' as a seed dressing at 1 g a.i. per kg of seed
BENOD F	Benodanil as a foliar spray at 1.1 kg
BTS S	'BTS 40542' as a seed dressing at 0.2 g a.i. per kg of seed
BTS F	'BTS 40542' as a foliar spray at 0.4 kg
CARBOX S	Carboxin as a seed dressing at 1.5 g per kg of seed
CARBOX F	Carboxin as a foliar spray at 1.1 kg
EL228 S	'EL 228' as a seed dressing at 0.2 g a.i. per kg of seed
EL228 F	'EL 228' as a foliar spray at 0.04 kg a.i.
H719 S	'H 719' as a seed dressing at 1.5 g a.i. per kg of seed
H719 F	'H 719' as a foliar spray at 1.1 kg a.i.
OM S	Organo-mercury as a seed dressing ('Agrosan GN' at 2.2 g per kg of seed)
PP296 F	'PP 296' as a foliar spray at 0.125 kg
TRIAD S	Triadimefon as a seed dressing at 0.5 g per kg of seed

NOTES: (1) All seed dressing treatments (except organo-mercury) were sown at 220 kg. All remaining treatments were sown at 190 kg.
(2) Foliar sprays were applied on 1 June, 1979 in 340 l.

Basal applications: Manures: (10:23:23) at 250 kg. 'Nitro-Chalk' at 460 kg.
Weedkillers: Paraquat at 0.42 kg ion in 220 l. Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 0.25 kg in 220 l).

Seed: Kador.

Cultivations, etc.:— Deep-tine cultivated twice: 31 Aug, 1978 and 1 Sept.
Heavy spring-tine cultivated: 14 Sept. NPK applied: 17 Oct. Paraquat applied: 18 Oct. Rolled and disc harrowed: 23 Oct. Seed sown: 24 Oct.
N applied: 19 Apr, 1979. 'Brittox' applied: 14 May. Combine harvested: 31 Aug. Previous cropping: Wheat 1977, barley 1978.

NOTE: Foot and root rots were assessed in spring and summer. Leaf diseases were assessed in late summer.

79/R/WW/5

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGCIDE	
NONE	4.77
BAS389 S	5.13
BENOD F	4.50
BTS S	5.65
BTS F	5.38
CARBOX S	5.16
CARBOX F	4.85
EL228 S	5.43
EL228 F	4.89
H719 S	4.68
H719 F	5.11
OM S	5.55
PP296 F	5.43
TRIAD S	5.68
MEAN	5.16

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGCIDE
-----	-----
SED	0.407

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	26	0.498	9.7

GRAIN MEAN DM% 86.1

PLOT AREA HARVESTED 0.00195