

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1979

[Full Table of Content](#)



79/R/B/11 Controlled Drop Application of Tridemorph - Barley

Rothamsted Research

Rothamsted Research (1980) *79/R/B/11 Controlled Drop Application of Tridemorph - Barley ; Yields Of The Field Experiments 1979*, pp 311 - 312 - DOI: <https://doi.org/10.23637/ERADOC-1-45>

79/R/B/11

SPRING BARLEY

CONTROLLED DROP APPLICATION OF TRIDEMORPH

Object: To compare controlled drop application with conventional spraying on the deposition of spray material, control of mildew and on the yield of spring barley - Claycroft.

Sponsors: F.T. Phillips, A.J. Arnold, P. Etheridge.

Design: 3 randomised blocks of 11 plots.

Whole plot dimensions: 4.27 x 24.4.

Treatments: All combinations of:-

- | | |
|-------------|--|
| 1. SPRAYER | Sprayer and drop density: |
| CDA 1 | Controlled drop application sprayer, standard drop density |
| CDA 2 | Controlled drop application sprayer, twice standard drop density |
| HYDRAUL | Hydraulic sprayer |
| 2. TRI RATE | Rates of applying tridemorph (on 12 June, 1979): |
| 1 | Standard, 525 g |
| 1/2 | Half standard, 263 g |
| 1/4 | Quarter standard, 132 g |
| EXTRA | plus two extra plots |
| NONE | Unsprayed |
| CDA R 1 | Controlled drop application sprayer, reduced drop density, applying standard rate tridemorph |

NOTES: (1) CDA sprayer applied tridemorph in 19 l.
(2) Hydraulic sprayer applied tridemorph in 340 l.

Basal applications: Manures: (20:14:14) at 440 kg, combine drilled. Weedkillers: Bromoxynil with ioxynil (as 'Oxytril CM' at 2.1 kg) and mecoprop at 1.6 kg in 220 l.

Seed: Wing, sown at 160 kg.

Cultivations, etc.: Subsoiled, tines 100 cm apart and 45 cm deep: 31 Oct, 1978.
Ploughed: 2 Nov. Spring-tine cultivated: 16 Apr, 1979. Seed sown: 17 Apr.
Weedkillers applied: 1 June. Combine harvested: 26 Aug. Previous crops: Beans 1977, wheat 1978.

NOTE: Observations were made on patterns of spray deposition using very small quantities of permethrin as a chemical marker.

79/R/B/11

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TRI RATE SPRAYER	1	1/2	1/4	MEAN
CDA 1	6.14	6.11	5.69	5.98
CDA 2	5.75	6.21	5.89	5.95
HYDRAUL	6.01	5.94	6.05	6.00
MEAN	5.96	6.09	5.88	5.98

EXTRA	NONE	CDA R 1	MEAN
	5.63	6.16	5.89

GRAND MEAN 5.96

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	SPRAYER	TRI RATE	SPRAYER TRI RATE & EXTRA
SED	0.144	0.144	0.249

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	20	0.305	5.1

GRAIN MEAN DM% 81.5

PLOT AREA HARVESTED 0.00520