

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/W/CS/253 Late N

Rothamsted Research

Rothamsted Research (1981) *80/W/CS/253 Late N* ; Yields Of The Field Experiments 1980, pp 194 - 196 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/W/CS/253

LATE N

Object: To study the effects of a range of fertilisers that release nitrogen later in the growing season than traditional forms on the growth and yield of potatoes - Woburn Lansome III.

Sponsors: F.V. Widdowson, A. Penny, T.M. Addiscott.

The first year, potatoes.

Design: 3 randomised blocks of 16 plots.

Whole plot dimensions: 4.27 x 13.1.

Treatments: All combinations of:-

- | | |
|-----------|---|
| 1. N FORM | Forms of nitrogen fertiliser: |
| AQ U | Aqueous urea, injected before planting |
| AQ U+CS2 | Aqueous urea + carbon disulphide at 10 kg, injected before planting |
| AQ U+NIT | Aqueous urea + nitrapyrin at 1 kg, injected before planting |
| NC E | 'Nitro-Chalk', all to the seedbed |
| NC E+L | 'Nitro-Chalk', half to the seedbed, half in June |
| IBDU | Isobutylidene diurea, all to the seedbed |

- | | |
|-----------|--------------------------------------|
| 2. N RATE | Rates of nitrogen fertiliser (kg N): |
|-----------|--------------------------------------|

200
300

plus four extra treatments all given 'Nitro-Chalk':

EXTRA

- | | |
|----------|--|
| NC E100 | At 100 kg N, all to the seedbed |
| NC E400 | At 400 kg N, all to the seedbed |
| NC EL100 | At 100 kg N, half to the seedbed, half in June |
| NC EL400 | At 400 kg N, half to the seedbed, half in June |

NOTE: Aqueous fertilisers were injected on 17 Apr, 1980. IBDU and seedbed 'Nitro-Chalk' were applied on 18 Apr. Late 'Nitro-Chalk' was applied on 10 June.

Basal applications: Manures: FYM at 50 t, (0:14:28) at 1880 kg. Weedkiller: Linuron at 1.1 kg in 280 l. Fungicide: Mancozeb at 1.3 kg in 300 l on five occasions with insecticide on the first, third, fourth and fifth occasions. Insecticide: Pirimicarb at 0.14 kg. Haulm desiccant: undiluted BOV at 170 l.

Seed: Pentland Crown.

80/W/CS/253

Cultivations, etc.:— Subsoiled, tines 160 cm apart and 40 cm deep: 12 Aug, 1979. FYM applied, ploughed: 21 Aug. PK applied: 12 Apr, 1980. Heavy spring-tine cultivated: 14 Apr. Spring-tine cultivated with crumbler attached: 17 Apr. Rotary cultivated, potatoes planted: 21 Apr. Weedkiller applied: 15 May. Earthed up: 12 June. Fungicide applied: 18 June, 3 July, 22 July, 8 Aug, 22 Aug. Insecticide applied: 18 June, 22 July, 8 Aug, 22 Aug. Haulm mechanically destroyed: 19 Sept. Haulm desiccant applied: 23 Sept. Lifted: 13 Oct. Previous crops: Barley 1978, grass ley 1979.

NOTE: Dry matter of tubers and haulm and numbers of tubers were assessed on several occasions during the season. Nitrogen contents of tubers were measured.

80/W/CS/253

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

N RATE	200	300	MEAN
N FORM			
AQ U	62.9	75.9	69.4
AQ U+CS2	65.4	67.3	66.3
AQ U+NIT	64.5	73.6	69.0
NC E	67.4	76.9	72.2
NC E+L	70.4	75.3	72.8
IBDU	71.1	73.1	72.1
MEAN	67.0	73.7	70.3

EXTRA	NC E100	NC E400	NC EL100	NC EL400	MEAN
	54.4	71.2	63.3	78.3	66.8

GRAND MEAN 69.4

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	N FORM	N RATE	N FORM N RATE & EXTRA
SED	3.53	2.50	1.44	3.53

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	30	4.32	6.2

PERCENTAGE WARE 4.44 CM (1.75 INCH) RIDDLE

***** TABLES OF MEANS *****

N RATE	200	300	MEAN
N FORM			
AQ U	91.1	94.9	93.0
AQ U+CS2	92.7	91.7	92.2
AQ U+NIT	92.8	94.0	93.4
NC E	92.1	92.7	92.4
NC E+L	94.2	93.9	94.0
IBDU	94.7	94.3	94.5
MEAN	92.9	93.6	93.3

EXTRA	NC E100	NC E400	NC EL100	NC EL400	MEAN
	90.0	92.3	92.8	95.0	92.5

GRAND MEAN 93.1

PLOT AREA HARVESTED 0.00260