

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/R/WW/4 Winter Wheat Seed and Divided N Dressings

Rothamsted Research

Rothamsted Research (1981) *80/R/WW/4 Winter Wheat Seed and Divided N Dressings* ; Yields Of The Field Experiments 1980, pp 229 - 230 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/R/WW/4

WINTER WHEAT

SEED RATES AND DIVIDED N DRESSINGS

Object: To study the effects of a range of rates of early nitrogen dressings on the growth and yield of w. wheat sown at half or standard seed rate - Gt. Knott III.

Sponsors: J. McEwen, R. Moffitt.

Design: 2 randomised blocks of 24 plots.

Whole plot dimensions: 4.27 x 8.08.

Treatments: All combinations of:-

1. SEEDRATE Seed rates (kg):

100
200

2. EARLY N Nitrogen fertiliser applied 4 Mar, 1980 (kg N):

0
25
50
75

3. APRIL N Nitrogen fertiliser applied 14 Apr (kg N):

75
100
125

Basal applications: Manures: (0:20:20) at 310 kg. Weedkillers: Paraquat at 0.56 kg ion in 220 l. Methabenzthiazuron at 1.5 kg in 220 l. Isoproturon at 2.9 kg in 250 l. Fungicide: Triadimefon at 0.13 kg in 250 l. Insecticide: Demeton-s-methyl at 0.24 kg in 250 l. Growth regulator: Chlormequat at 1.7 l in 250 l.

Seed: Flanders.

Cultivations, etc.: Heavy spring-tine cultivated twice: 24 Sept, 1979. Paraquat applied: 15 Oct. PK applied, spring-tine cultivated, seed sown: 22 Oct. Methabenzthiazuron applied: 24 Oct. Isoproturon applied: 24 Apr, 1980. Growth regulator applied: 7 May. Fungicide applied: 3 June. Insecticide applied: 23 June. Combine harvested: 22 Aug. Previous crops: S. barley 1978, s. oats 1979.

NOTES: (1) Plant counts were made in January, tiller counts in April and ear counts in July.

(2) 1000 grain weights and N content of grain were measured.

80/R/WW/4

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

EARLY N SEEDRATE	0	25	50	75	MEAN
100	6.66	7.32	7.94	7.83	7.44
200	7.22	7.36	7.80	7.86	7.56
MEAN	6.94	7.34	7.87	7.85	7.50
APRIL N SEEDRATE	75	100	125	MEAN	
100	7.29	7.38	7.64	7.44	
200	7.13	7.68	7.87	7.56	
MEAN	7.21	7.53	7.76	7.50	
APRIL N EARLY N	75	100	125	MEAN	
0	6.56	6.96	7.31	6.94	
25	6.87	7.46	7.69	7.34	
50	7.65	8.10	7.85	7.87	
75	7.77	7.59	8.18	7.85	
MEAN	7.21	7.53	7.76	7.50	
SEEDRATE	APRIL N EARLY N	75	100	125	
100	0	6.22	6.67	7.10	
	25	7.12	7.36	7.47	
	50	8.05	7.94	7.84	
	75	7.77	7.56	8.16	
200	0	6.90	7.25	7.51	
	25	6.62	7.57	7.90	
	50	7.25	8.27	7.86	
	75	7.77	7.62	8.20	

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	SEEDRATE	EARLY N	APRIL N	SEEDRATE EARLY N
SED	0.089	0.126	0.109	0.178
TABLE	SEEDRATE APRIL N	EARLY N APRIL N	SEEDRATE EARLY N APRIL N	
SED	0.154	0.218	0.308	

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	23	0.308	4.1

GRAIN MEAN DM% 84.0 PLOT AREA HARVESTED 0.00246