

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/R/WS/1 Spring Wheat Fungicides and Alternaria

Rothamsted Research

Rothamsted Research (1981) *80/R/WS/1 Spring Wheat Fungicides and Alternaria* ; Yields Of The Field Experiments 1980, pp 248 - 250 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/R/WS/1

SPRING WHEAT

FUNGICIDES AND ALTERNARIA

Object: To study the effects of a range of fungicides, and times of application, on the incidence of fungi, especially *Alternaria*, on the ripening grain and on the yield of s. wheat - Delafield.

Sponsor: N. Magan.

Design: 2 randomised blocks of 24 plots.

Whole plot dimensions: 4.27 x 16.2.

Treatments: All combinations of:-

1. E FUNG Early-applied fungicide:

NONE	None
CARB+MAN	Carbendazim at 0.25 kg plus maneb at 1.6 kg applied on 9 June 1980

2. L FUNG Late-applied fungicide:

BENOMYL	Benomyl at 0.56 kg
PROCHLOR	Prochloraz at 0.50 kg
CAPTAFOL	Captafol at 1.40 kg
CARB+MAN	Carbendazim at 0.25 kg plus maneb at 1.6 kg
IMAZALIL	Imazalil at 0.5 kg

3. LFNGDATE Dates of applying late fungicide:

25 JUNE
7 JULY

plus two extra treatments not given L FUNG:

L FNG 0

NONE	No early-applied fungicide (duplicated)
CARB+MAN	Carbendazim at 0.25 kg plus maneb at 1.6 kg applied on 9 June (duplicated)

NOTE: Treatment sprays were applied in 340 l.

Basal applications: Manures: (20:10:10) at 450 kg, combine drilled.
Weedkillers: Dicamba with mecoprop and MCPA (as 'Banlene Plus' at 5.0 l) in 250 l.

Seed: Timmo, sown at 190 kg.

Cultivations, etc.: Subsoiled with tines 160 cm apart and 40 cm deep: 2 Nov, 1979. Chisel ploughed twice: 4 Dec. Spring-tine cultivated, seed sown: 6 Apr, 1980. Weedkillers applied: 27 May. Combine harvested: 4 Sept. Previous crops: S. beans and kale 1978, potatoes 1979.

80/R/WS/1

NOTES: Grain microflora, especially *Alternaria*, were assessed at fortnightly intervals after heading. Thousand grain weights were measured, and grain was assessed for germination and seedling growth.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

L FUNG	BENOMYL	PROCHLOR	CAPTAFOL	CARB+MAN	IMAZALIL	MEAN
E FUNG						
NONE	4.33	4.51	4.51	4.50	4.14	4.40
CARB+MAN	4.86	4.75	4.72	4.50	4.83	4.73
MEAN	4.59	4.63	4.62	4.50	4.48	4.56
LFNGDATE	25 JUNE	7 JULY	MEAN			
E FUNG						
NONE	4.39	4.40	4.40			
CARB+MAN	4.78	4.68	4.73			
MEAN	4.59	4.54	4.56			
LFNGDATE	25 JUNE	7 JULY	MEAN			
L FUNG						
BENOMYL	4.77	4.41	4.59			
PROCHLOR	4.56	4.70	4.63			
CAPTAFOL	4.60	4.64	4.62			
CARB+MAN	4.56	4.43	4.50			
IMAZALIL	4.44	4.53	4.48			
MEAN	4.59	4.54	4.56			
E FUNG	LFNGDATE	25 JUNE	7 JULY			
NONE	L FUNG					
	BENOMYL	4.37	4.28			
	PROCHLOR	4.32	4.70			
	CAPTAFOL	4.50	4.52			
	CARB+MAN	4.56	4.44			
	IMAZALIL	4.22	4.07			
CARB+MAN	BENOMYL	5.18	4.54			
	PROCHLOR	4.80	4.70			
	CAPTAFOL	4.69	4.75			
	CARB+MAN	4.57	4.42			
	IMAZALIL	4.66	5.00			
L FUNG 0	NONE	CARB+MAN	MEAN			
	4.44	4.92	4.68			

GRAND MEAN 4.58

80/R/WS/1

GRAIN TONNES/HECTARE

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	L FNG 0	E FUNG	L FUNG	LFNGDATE
SED	0.285	0.127	0.202	0.127

TABLE	E FUNG L FUNG	E FUNG LFNGDATE	L FUNG LFNGDATE	E FUNG L FUNG LFNGDATE
SED	0.285	0.180	0.285	0.403

SED FOR COMPARING A MEAN IN E FUNG.L FUNG.LFNGDATE TABLE WITH A L FNG 0
MEAN IS 0.247

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	25	0.403	8.8

GRAIN MEAN DM% 79.1

PLOT AREA HARVESTED 0.00260