

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/R/AG/6 Agdell

Rothamsted Research

Rothamsted Research (1981) *80/R/AG/6 Agdell* ; Yields Of The Field Experiments 1980, pp 26 - 29 -
DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/R/AG/6

AGDELL

Object: To study, by crop yields and soil analyses, the residual values of phosphate and potash applied in the period 1848-1951 and further dressings since 1964.

The eleventh year of revised scheme, s. beans, w. wheat.

For previous years see 'Details' 1967 and 1973, and 74-79/R/AG/6.

Treatments: All combinations of:-

Whole plots

1. OLDRESD Fertilisers and organic manures applied to roots every fourth year, in the period 1848-1948:

NONE	None
PKNAMG	P K Na Mg
NPKNAMGC	N P K Na Mg C

N: 48 kg N as sulphate of ammonia
P: 41 kg P as superphosphate
K: 224 kg K as sulphate of potash
Na: 16 kg Na as sulphate of soda
Mg: 11 kg Mg as sulphate of magnesia
C: Castor meal at 2240 kg supplying about 112 kg N

2. RN CROP Rotation 1848-1951 and crop in 1980:

F/WHEAT	With fallow: Roots (turnips or swedes), s. barley, fallow, w. wheat 1848-1951. W. wheat (after s. beans 1979)
L/BEANS	With legume: Roots, s. barley, legume (clover or beans), w. wheat 1848-1951. S. beans (after grass 1979)

Half plots

3. 1964RESD Residues of 1964 treatments:

P
K

Quarter plots

4. PREVCROP Previous cropping 1958-69 on P-test half plots, 1958-70 on K-test half plots:

ARABLE	Arable or fallow
GRASS	Grass

80/R/AG/6

Sixteenth plots

5. P₂O₅ 64 K₂O 64

Rates of 1964 treatments (kg):
P₂O₅ to P-test K₂O to K-test
half plots half plots

0	0
500	315
1000	630
2000	1260

Thirty second plots

6.

On P-test half plots:
To RN CROP F/WHEAT. Residues of P₂O₅ applied
1970-72 (total, kg) and 1979 (kg):

P₂O₅ 729

(0)0
(375)150

To RN CROP L/BEANS. Residues of P₂O₅ applied
1970-72 (total, kg) and a fresh dressing in 1980
(kg):

P₂O₅ 720

(0)0
(375)150

On K-test half plots:
To RN CROP F/WHEAT. Residues of K₂O applied
1973-76 (total, kg) and 1979 (kg):

K₂O 769

(0)0
(870)300

To RN CROP L/BEANS. Residues of K₂O applied
1973-76 (total, kg) and a fresh dressing in 1980
(kg):

K₂O 760

(0)0
(870)300

Standard applications:

W. wheat: Manures: 'Nitro-Chalk' at 580 kg. Weedkillers: Glyphosate at 1.5 kg in 220 l. Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 3.5 l) in 900 l.

W. beans: Weedkillers: Trietazine with simazine (as 'Remtal SC' at 2.5 l) in 250 l. Permethrin at 0.13 kg in 900 l. Pirimicarb at 0.14 kg in 250 l.

80/R/AG/6

Seed: Wheat: Flanders, sown at 200 kg.

Beans: Minden, sown at 220 kg.

Cultivations, etc.:-

Wheat: Glyphosate applied: 27 Sept, 1979. Disc harrowed twice, seed sown: 15 Oct. N applied: 15 Apr, 1980. 'Brittox' applied: 25 Apr. Combine harvested: 21 Aug.

Beans: Test P and K applied: 17 July, 1979. Ploughed: 18 July. Heavy spring-tine cultivated: 31 July, 10 Aug. Rotary harrowed: 4 Mar, 1980. Seed sown: 5 Mar. Weedkiller applied: 21 Mar. Permethrin applied: 6 May. Pirimicarb applied: 16 July. Combine harvested: 18 Sept.

SPRING BEANS P PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	OLDRES	NONE	PKNAMG	NPKNAMG
	P205 720	(0)0 (375)150	(0)0 (375)150	(0)0 (375)150
PREVCROP	P205 64			
ARABLE	0	2.13	2.52	1.41
	500	1.50	1.42	1.34
	1000	1.43	1.49	2.77
	2000	2.30	1.82	2.12
GRASS	0	1.67	2.21	1.79
	500	2.55	3.11	1.55
	1000	1.59	1.74	2.06
	2000	2.62	3.12	2.57

GRAIN MEAN DM% 78.9

SPRING BEANS K PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	OLDRES	NONE	PKNAMG	NPKNAMG
	K20 760	(0)0 (870)300	(0)0 (870)300	(0)0 (870)300
PREVCROP	K20 64			
ARABLE	0	0.94	2.68	1.39
	315	1.00	1.43	1.35
	630	1.01	1.27	1.03
	1260	1.90	2.60	1.41
GRASS	0	1.01	1.85	1.12
	315	1.78	2.96	1.34
	630	1.20	2.82	1.42
	1260	0.93	2.23	1.84

GRAIN MEAN DM% 78.6

PLOT AREA HARVESTED (OLDRES NONE) 0.00117

PLOT AREA HARVESTED (REMAINDER) 0.00132

80/R/AG/6

WINTER WHEAT P PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	OLDRESO	NONE		PKNAMG		NPKNAMGC	
	P205 729	(0)0	(375)150	(0)0	(375)150	(0)0	(375)150
PREVCROP	P205 64						
ARABLE	0	4.28	5.20	4.21	4.66	5.64	6.26
	500	4.26	5.07	5.36	5.61	6.56	5.50
	1000	5.30	5.54	4.64	5.30	4.94	5.74
	2000	4.54	4.80	6.04	6.20	6.29	6.64
GRASS	0	4.54	5.85	3.48	4.89	5.12	5.40
	500	4.46	4.16	5.89	6.25	4.67	4.94
	1000	4.87	5.43	5.43	5.50	5.11	6.60
	2000	5.07	5.73	6.39	6.36	4.88	5.33

GRAIN MEAN DM% 82.3

WINTER WHEAT K PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	OLDRESO	NONE		PKNAMG		NPKNAMGC	
	K20 769	(0)0	(870)300	(0)0	(870)300	(0)0	(870)300
PREVCROP	K20 64						
ARABLE	0	5.59	4.27	5.80	5.13	5.37	6.03
	315	4.73	4.56	5.22	5.76	5.73	6.16
	630	4.60	4.97	4.57	4.84	5.89	5.73
	1260	4.79	4.97	5.09	5.22	6.00	5.70
GRASS	0	4.58	4.42	5.73	5.43	6.07	5.99
	315	5.13	5.15	5.92	5.99	6.15	6.35
	630	4.99	5.89	5.74	5.71	4.54	6.05
	1260	5.52	5.52	5.99	5.92	5.69	6.22

GRAIN MEAN DM% 82.4

PLOT AREA HARVESTED (OLDRESO NONE) 0.00120

PLOT AREA HARVESTED (REMAINDER) 0.00134