

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/W/P/5 Potatoes Varieties and Potato Cyst Nematode

Rothamsted Research

Rothamsted Research (1981) *80/W/P/5 Potatoes Varieties and Potato Cyst Nematode* ; Yields Of The Field Experiments 1980, pp 329 - 330 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/W/P/5

POTATOES

VARIETIES AND POTATO CYST NEMATODE

Object: To measure the yields of potato varieties, resistant and susceptible to potato cyst-nematode, on infested land treated with and without oxamyl and to study effects on nematode populations - Woburn, Horsepool.

Sponsor: A.G. Whitehead.

Design: 3 randomised blocks of 16 plots.

Whole plot dimensions: 2.84 x 6.09.

Treatments: All combinations of:-

1. VARIETY	Varieties:
A BANNER	Arran Banner
CARA	Cara
CROFT	Croft
DESIREE	Desiree
M PIPER	Maris Piper
P CROWN	Pentland Crown
P DELL	Pentland Dell
RECORD	Record

2 OXAMYL	Oxamyl to seedbed (kg):
0.0	
5.6	

Basal applications:- Manures: (13:13:20) at 1880 kg. Weedkiller: Linuron at 1.1 l in 280 l. Fungicide: Mancozeb at 1.3 kg in 300 l applied five times, with insecticide on the third, fourth and fifth occasions. Insecticide: Pirimicarb at 0.14 kg. Haulm desiccant: Undiluted BOV at 170 l.

Cultivations, etc.:- Heavy spring-tine cultivated: 20 Oct, 1979. Spring-tine cultivated: 22 Oct. NPK applied: 12 Apr, 1980. Heavy spring-tine cultivated: 14 Apr. Oxamyl applied, rotary cultivated, potatoes planted: 1 May. Weedkiller applied: 16 May. Fungicide applied: 18 June, 3 July, 22 July, 10 Aug, 22 Aug. Insecticide applied: 22 July, 10 Aug, 22 Aug. Haulm desiccant applied: 24 Sept. Lifted: 30 Sept. Previous crops: W. oats 1978, potatoes 1979.

NOTES: (1) Soil samples were taken before applying treatments and after harvest for counts of cysts, eggs, and larvae of *Globodera rostochiensis*.

(2) The treatment combinations of OXAMYL 0.0 with VARIETY A BANNER, CROFT, DESIREE, P CROWN, P DELL, RECORD gave very poor yields. These combinations were omitted from the analysis presented and standard errors presented do not apply to them.

80/W/P/5

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

OXAMYL	0.0	5.6
VARIETY		
A BANNER	0.3	53.0
CARA	37.1	63.4
CROFT	3.2	65.7
DESIREE	1.3	52.1
M PIPER	24.8	63.7
P CROWN	2.5	65.7
P DELL	1.0	52.6
RECORD	1.1	61.4

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY*
	OXAMYL
-----	-----
SED	5.10

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE*	CV%
BLOCK.WP	18	6.25	11.6

* CALCULATED ONLY FROM PLOTS WITH VARIETIES
CARA AND M PIPER WITH OXAMYL 0.0 AND
ALL VARIETIES WITH OXAMYL 5.6

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE TONNES/HECTARE

***** TABLES OF MEANS *****

OXAMYL	0.0	5.6
VARIETY		
A BANNER	0.0	97.3
CARA	96.5	97.8
CROFT	52.4	95.9
DESIREE	0.0	93.4
M PIPER	94.1	94.4
P CROWN	34.8	96.4
P DELL	9.5	96.6
RECORD	7.0	96.0

PLOT AREA HARVESTED 0.00087