

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/W/CS/253 Late N - W. Wheat

Rothamsted Research

Rothamsted Research (1982) *81/W/CS/253 Late N - W. Wheat* ; Yields Of The Field Experiments 1981, pp 193 - 194 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/W/CS/253

LATE N

Object: To study the residual effects on w. wheat of a range of fertilisers applied to potatoes in 1980 - Woburn Lansome III.

Sponsors: F.V. Widdowson, A. Penny, T.M. Addiscott.

The second year, w. wheat.

For previous year see 80/W/CS/253.

Design: 3 randomised blocks of 16 plots.

Whole plot dimensions: 4.27 x 9.1.

Treatments: All combinations of:-

1. N FORM (80) Forms of nitrogen fertiliser in 1980:

AQ U	Aqueous urea, injected before planting
AQ U+CS2	Aqueous urea + carbon disulphide at 10 kg, injected before planting
AQ U+NIT	Aqueous urea + nitrapyrin at 1 kg, injected before planting
NC E	'Nitro-Chalk', all to the seedbed
NC E+L	'Nitro-Chalk', half to the seedbed, half in June
IBDU	Isobutylidene diurea, all to the seedbed
2. N RATE (80) Rates of nitrogen fertiliser in 1980 (kg N):

200
300

plus four extra treatments all given 'Nitro-Chalk' in 1980:

EXTRA

- | | |
|----------|--|
| NC E100 | At 100 kg N, all to the seedbed |
| NC E400 | At 400 kg N, all to the seedbed |
| NC EL100 | At 100 kg N, half to the seedbed, half in June |
| NC EL400 | At 400 kg N, half to the seedbed, half in June |

Basal applications: Manures: (0:20:20) at 310 kg. N at 120 kg as 'Nitro-Chalk'. Weedkillers: Isoproturon at 2.4 l in 280 l. Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 2.5 l) in 280 l. Fungicides: Prochloraz at 0.4 l with maneb at 1.2 kg and zineb at 0.13 kg in 280 l. Growth regulator: Chlormequat at 1.4 l in 280 l.

Seed: Flanders, sown at 190 kg.

Cultivations, etc.: Heavy spring-tine cultivated: 15 Oct, 1980, 25 Oct.
PK applied: 29 Oct. Seed sown, spring-tine cultivated in: 30 Oct.
N applied: 9 Apr, 1981. Growth regulator applied: 20 Apr.
Weedkillers applied: 22 Apr. Fungicides applied: 18 June. Combine harvested: 24 Aug.

81/W/CS/253

GRAIN DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

N RATE(80)	200	300	MEAN
N FORM(80)			
AQ U	7.35	7.86	7.61
AQ U+CS2	7.79	7.86	7.83
AQ U+NIT	7.16	8.06	7.61
NC E	7.59	8.06	7.82
NC E+L	7.85	7.71	7.78
IBDU	7.56	7.51	7.54
MEAN	7.55	7.84	7.70

EXTRA	NC E100	NC E400	NC EL100	NC EL400	MEAN
	7.26	7.75	7.86	7.26	7.53

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	N FORM(80)	N RATE(80)	N FORM(80) N RATE(80) & EXTRA
SED	0.458	0.324	0.187	0.458

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	30	0.560	7.3

GRAIN MEAN DM% 82.5

PLOT AREA HARVESTED 0.00279