

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/R/CS/264 Fungicide Rates - W. Oilseed Rape, S. Barley

Rothamsted Research

Rothamsted Research (1982) *81/R/CS/264 Fungicide Rates - W. Oilseed Rape, S. Barley* ; Yields Of The Field Experiments 1981, pp 209 - 211 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/CS/264

FUNGICIDE RATES

Object: To study the effects of a range of rates of triadimefon on pathogens and yield of w. oilseed rape on land in rotation (pathogen free) and after previous oilseed rape (pathogen infected) - Long Hoos IV 1 (pathogen free) and Summerdells I (pathogen infected).

Sponsor: C.J. Rawlinson.

The first year, w. oilseed rape, s. barley.

Design: 3 randomised blocks of 6 plots.

Whole plot dimensions: 2.13 x 3.05.

Treatments:

FUNGRATE	Rates of triadimefon (kg) applied to soil surface on 28 Aug, 1980 before seedbed cultivation:
----------	---

0.06
0.12
0.25
0.50
1.00
2.00

NOTES: (1) Sprays were applied in 340 l.

(2) On Summerdells I w. oilseed rape failed during the winter and the site was sown to s. barley to assess the residual effects of treatments already applied.

Standard applications: Long Hoos IV and Summerdells I: Manures: 'Nitro-Chalk' at 670 kg. Weedkillers: Propyzamide at 0.56 kg with dalapon at 1.9 kg in 340 l. Manures: (13:13:20) at 380 kg. Weedkillers: Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 3 l.) in 250 l.

Seed: Long Hoos IV and Summerdells I: Primor, sown at 9 kg. Summerdells I: Georgie, sown at 160 kg.

Cultivations, etc.:-

Long Hoos IV: NPK applied: 1 Sept, 1980. Rotary harrowed, seed sown: 5 Sept. Weedkillers applied: 30 Oct. N applied: 18 Feb, 1981. Cut and windrowed: 21 July. Grain threshed: 31 July. Previous crops: S. barley 1979, fallow 1980.

Summerdells I: NPK applied: 1 Sept, 1980. Heavy spring-tine cultivated twice: 2 Sept. Rotary harrowed: 4 Sept. Primor seed sown: 5 Sept. Propyzamide and dalapon applied: 30 Oct. N applied: 18 Feb, 1981. Chisel ploughed: 1 May. Rotary harrowed: 13 May. Georgie seed sown: 14 May. 'Brittox' applied: 19 June. Combine harvested: 8 Sept. Previous crops: W. barley 1979, w. oilseed rape 1980.

81/R/CS/264

- NOTES: (1) On Summerdells I: Foliar diseases on w. oilseed rape were assessed twice; s. barley was assessed fortnightly for mildew, yellow and brown rust. Soil cores were analysed for triadimefon residues.
- (2) On Long Hoos IV: Foliar diseases were assessed five times during the season.
- (3) 2 plots, with FUNGRATE 0.06 and 2.00 were lost, estimated values were used in the analysis.

81/R/CS/264 LONG HOOS IV

OILSEED RAPE

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGRATE	0.06	0.12	0.25	0.50	1.00	2.00	MEAN
	1.41	1.24	1.24	1.44	1.29	1.28	1.32

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGRATE
-----	-----
SED	0.079

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	10	0.079	7.4

GRAIN MEAN DM% 78.4

PLOT AREA HARVESTED 0.00065

81/R/CS/264 SUMMERDELLS I

SPRING BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGRATE	0.06	0.12	0.25	0.50	1.00	2.00	MEAN
	2.35	1.92	2.22	2.48	2.49	2.82	2.38

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGRATE
-----	-----
SED	0.257

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	8	0.314	13.2

GRAIN MEAN DM% 79.4

PLOT AREA HARVESTED 0.00053