

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/R/CS/265 Soil Fumigation, Mycorrhiza and P - S. Wheat

Rothamsted Research

Rothamsted Research (1982) *81/R/CS/265 Soil Fumigation, Mycorrhiza and P - S. Wheat* ; Yields Of The Field Experiments 1981, pp 212 - 214 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/CS/265

SOIL FUMIGATION, MYCORRHIZA AND P

Object: To study the effects on s. wheat of applications of mycorrhizal inoculum, methyl bromide and rates of phosphate fertiliser - Delharding.

Sponsors: J.A. Buwalda, D.P. Stribley, P.B. Tinker.

The first year, s. wheat.

Design: 3 randomised blocks of 8 plots split into 2.

Whole plot dimensions: 4.4 x 3.0.

Treatments: All combinations of:-

Whole plots

1. STERILNT Soil sterilant:

NONE	None
METH BR	Methyl bromide at 980 kg

2. P Rates of phosphate fertiliser (kg P), as superphosphate:

0
15
30
60

Sub plots

3. INOCULUM Mycorrhizal inoculum:

NONE	None
G MOSSE	Glomus mosseae

NOTE: Inoculum was prepared by growing leeks in pots of soil infected with the mycorrhiza. After 20 weeks growth, soil and roots in the pots were chopped and applied to the seed furrows at 3.5 t per ha. Uninoculated plots received soil and roots at the same rate from pots growing uninfected leeks.

Basal applications: Manures: Dolomitic limestone at 12.5 t. Chalk at 3.0 t N at 125 kg as 'Nitro-Chalk', K at 50 kg as muriate of potash, Mg at 100 kg as Epsom salts. Weedkillers: Ioxynil at 0.42 l and mecoprop at 1.26 l in 250 l, applied with the fungicide. Fungicide: Tridemorph at 0.53 kg.

Seed: Highbury, sown at 350 seeds per m² (about 170 kg).

Cultivations, etc.: - Dolomitic limestone applied: 15 Sept, 1980. Epsom salts applied: 13 Jan, 1981. Basal chalk, N, and K and test P applied: 18 Mar. Methyl bromide applied: 13 Apr. Inoculum applied, seed sown: 23 Apr. Weedkillers applied: 7 June. Harvested by hand: 9 Sept.

NOTE: Plots were sampled five times during the season to assess mycorrhizal infection, and twice to measure P content of the leaves and soil.

81/R/CS/265

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

P	0	15	30	60	MEAN
STERILNT					
NONE	2.70	3.86	4.31	5.07	3.98
METH BR	2.70	3.62	4.23	4.90	3.86
MEAN	2.70	3.74	4.27	4.98	3.92
INOCULUM	NONE	G MOSSE	MEAN		
STERILNT					
NONE	3.86	4.11	3.98		
METH BR	3.39	4.33	3.86		
MEAN	3.62	4.22	3.92		
INOCULUM	NONE	G MOSSE	MEAN		
P					
0	2.31	3.09	2.70		
15	3.45	4.03	3.74		
30	4.00	4.54	4.27		
60	4.73	5.23	4.98		
MEAN	3.62	4.22	3.92		
STERILNT	INOCULUM	NONE	G MOSSE		
NONE	P				
	0	2.42	2.99		
	15	3.84	3.89		
	30	4.16	4.45		
	60	5.01	5.12		
METH BR	0	2.21	3.19		
	15	3.07	4.16		
	30	3.83	4.64		
	60	4.45	5.34		

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	STERILNT	P	INOCULUM	STERILNT P
SED	0.201	0.285	0.075	0.403
TABLE	STERILNT INOCULUM	P INOCULUM	STERILNT P INOCULUM	
SED	0.215	0.304	0.429	
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:				
STERILNT	0.106			
P		0.149		
STERILNT.P			0.211	

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.STERILNT.P	14	0.493	12.6
BLOCK.STERILNT.P.INOCULUM	16	0.258	6.6

81/R/CS/265

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

P	0	15	30	60	MEAN
STERILNT					
NONE	3.38	4.40	5.01	5.67	4.61
METH BR	4.47	5.31	5.83	5.80	5.35
MEAN	3.93	4.85	5.42	5.73	4.98

INOCULUM	NONE	G MOSSE	MEAN
STERILNT			
NONE	4.41	4.82	4.61
METH BR	4.95	5.76	5.35
MEAN	4.68	5.29	4.98

INOCULUM	NONE	G MOSSE	MEAN
P			
0	3.44	4.41	3.93
15	4.49	5.22	4.85
30	5.11	5.73	5.42
60	5.67	5.80	5.73
MEAN	4.68	5.29	4.98

	INOCULUM	NONE	G MOSSE
STERILNT	P		
NONE	0	3.02	3.74
	15	4.30	4.50
	30	4.84	5.18
	60	5.48	5.86
METH BR	0	3.87	5.08
	15	4.68	5.94
	30	5.39	6.28
	60	5.86	5.73

NO DRY MATTER PERCENTAGES - CROP WEIGHED DRY

SUB PLOT AREA HARVESTED 0.00020