

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/R/RN/17 Rates of P and K to the Subsoil - Potatoes, S. Barley, S. Beans, W. Wheat

Rothamsted Research

Rothamsted Research (1982) *81/R/RN/17 Rates of P and K to the Subsoil - Potatoes, S. Barley, S. Beans, W. Wheat* ; Yields Of The Field Experiments 1981, pp 95 - 101 - DOI:
<https://doi.org/10.23637/ERADOC-1-35>

81/R/RN/17

RATES OF P AND K TO THE SUBSOIL

Object: To study the effects of a range of rates and frequencies of application of P & K to the subsoil, singly and together, on the yields and nutrient uptakes of a rotation of crops - Meadow.

Sponsors: J. McEwen, A.E. Johnston.

The first year, potatoes, s. barley, s. beans, w. wheat.

Design: 4 series (for crops) each of 40 plots.

Whole plot dimensions: 3.0 x 14.

Treatments to each series:

TREATMNT Extra P & K and primary cultivation tool:

	P ₂ O ₅ (kg)	K ₂ O(kg)	Tool	
- - -	0	0	Plough	(duplicated)
P6 K6 T	1000	500 to topsoil	"	(")
- - S	0	0 " "	Wye double-digger	(six plots)
P2 - S	63	0 to subsoil	" "	"
P3 - S	125	0 " "	" "	"
P4 - S	250	0 " "	" "	"
P5 - S	500	0 " "	" "	" (duplicated)
P6 - S	1000	0 " "	" "	"
- K2 S	0	31 " "	" "	"
- K3 S	0	63 " "	" "	"
- K4 S	0	125 " "	" "	"
- K5 S	0	250 " "	" "	" (duplicated)
- K6 S	0	350 " "	" "	"
P1 K1 S	31	16 " "	" "	"
P1 K3 S	31	63 " "	" "	"
P2 K2 S	63	31 " "	" "	"
P3 K1 S	125	16 " "	" "	"
P3 K3 S	125	63 " "	" "	"
P3 K4 S	125	125 " "	" "	"
P4 K3 S	250	63 " "	" "	"
P4 K4 S	250	125 " "	" "	"
P4 K5 S	250	250 " "	" "	" (duplicated)
P4 K6 S	250	350 " "	" "	"
P5 K4 S	500	125 " "	" "	" (duplicated)
P5 K5 S	500	250 " "	" "	"
P5 K6 S	500	350 " "	" "	"
P6 K4 S	1000	125 " "	" "	"
P6 K5 S	1000	250 " "	" "	"
P6 K6 S	1000	350 " "	" "	"

- NOTES: (1) Subsoiling was done with the Wye double-digger which turns a furrow with a conventional plough share, to a depth of 23 cm, and at the same time rotary cultivates the bottom of the adjacent furrow to a further depth of 15 cm. When applying P & K this was distributed ahead of the rotary cultivator.
- (2) The topsoil PK dressing was equally divided before and after ploughing.

81/R/RN/17

(3) Treatments shown are those for the first year only and provide for comparisons of frequency of application.

(4) The rate of 350 kg K₂O applied was in error for 500 kg K₂O.

Standard applications:

All crops: Weedkillers: Glyphosate at 1.5 l in 250 l.

Potatoes: Manures: (10:10:15 + 4.5 Mg) at 1960 kg. Weedkillers: Linuron at 1.13 with paraquat at 0.4 kg ion in 250 l. Fungicides: Mancozeb at 1.4 kg in 250 l applied six times, with pirimicarb on the first five. Insecticide: Pirimicarb at 0.14 kg.

S. barley: Manures: (20:10:10) at 450 kg. Weedkillers: Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 2.5 l) in 250 l applied with the fungicide. Fungicide: Tridemorph at 0.53 kg. Prochloraz at 0.4 l in 250 l.

S. beans: Weedkiller: Simazine at 1.2 kg in 250 l. Fungicide: Benomyl at 0.55 kg in 250 l applied with the pirimicarb. Insecticides: Phorate at 5.6 kg. Pirimicarb at 0.14 kg.

W. wheat: Manures: (0:20:20) at 310 kg, 'Nitro-Chalk' at 560 kg. Weedkillers: Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 3.7 l) in 250 l. Fungicides: Prochloraz at 0.4 l in 250 l applied twice, with maneb at 1.2 kg and zineb at 0.13 kg on the second occasion.

Seed: Potatoes: Pentland Crown.

S. barley: Triumph, seed dressed with ethirimol, sown at 160 kg.

S. beans: Minden, sown at 280 kg.

W. wheat: Flanders, sown at 200 kg.

Cultivations, etc.:-

All crops: Glyphosate applied: 24 Sept, 1980. Treatments applied by double digger: 23-31 Oct. Half topsoil PK treatment applied & ploughed in: 10 Nov. Second half topsoil PK treatment applied: 11 Nov.

Potatoes: Chisel ploughed: 21 Nov, 1980. NPK applied: 14 Apr, 1981.

Spike rotary cultivated, potatoes planted: 30 Apr. Weedkillers applied: 21 May. Fungicide with insecticide applied: 22 June, 1 July, 13 July, 27 July, and 12 Aug. Fungicide applied: 24 Aug. Haulm mechanically destroyed: 7 Oct. Lifted: 22 Oct.

S. barley: Chisel ploughed: 21 Nov, 1980. Spring-tine cultivated, NPK applied: 17 Feb, 1981. Spring-tine cultivated, seed sown by combined drill and rotary harrow: 18 Feb. Weedkillers and tridemorph applied: 12 May. Prochloraz applied: 8 July. Combine harvested: 17 Aug.

S. beans: Chisel ploughed: 21 Nov, 1980. Spring-tine cultivated: 17 Feb, 1981. Phorate applied, spring-tine cultivated, seed sown by combined drill and rotary harrow: 18 Feb. Weedkiller applied: 9 Apr. Pirimicarb and fungicide applied: 18 June. Combine harvested: 8 Sept.

W. wheat: Heavy spring-tine cultivated, PK applied, spring-tine cultivated: 11 Nov, 1980. Seed sown by combined drill and rotary harrow: 12 Nov. N applied: 13 Apr, 1981. Weedkillers applied: 22 Apr. Prochloraz applied: 5 May. Prochloraz applied with maneb and zineb: 19 June. Combine harvested: 24 Aug.

Previous crops: W. wheat 1979, s. barley 1980.

81/R/RN/17

POTATOES SERIES I

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	61.7
P6 K6 T	68.3
- - S	61.3
P2 - S	71.8
P3 - S	62.3
P4 - S	63.3
P5 - S	59.2
P6 - S	67.1
- K2 S	65.9
- K3 S	64.1
- K4 S	70.4
- K5 S	69.7
- K6 S	72.1
P1 K1 S	66.7
P1 K3 S	65.0
P2 K2 S	51.4
P3 K1 S	61.6
P3 K3 S	63.4
P3 K4 S	64.4
P4 K3 S	54.8
P4 K4 S	54.8
P4 K5 S	64.4
P4 K6 S	75.7
P5 K4 S	63.6
P5 K5 S	65.0
P5 K6 S	71.0
P6 K4 S	67.0
P6 K5 S	75.9
P6 K6 S	72.3
MEAN	64.7

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
SED	4.58 MIN REP
	3.74 MAX-MIN

* SED APPLIES ONLY TO - - -, P6 K6 T, - - S, P5 - S, - K5 S, P4 K6 S and P5 K4 S.

TREATMNT	
MAX-MIN	- - S V ANY OF REMAINDER
MIN REP	ANY OF REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	11	4.58	7.1

81/R/RN/17

POTATOES SERIES I

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

TREATMNT	
- - -	96.1
P6 K6 T	97.5
- - S	97.4
P2 - S	98.0
P3 - S	97.3
P4 - S	96.9
P5 - S	97.7
P6 - S	97.7
- K2 S	97.3
- K3 S	97.2
- K4 S	98.2
- K5 S	96.7
- K6 S	97.5
P1 K1 S	97.7
P1 K3 S	97.4
P2 K2 S	97.2
P3 K1 S	96.3
P3 K3 S	97.9
P3 K4 S	97.7
P4 K3 S	97.5
P4 K4 S	98.3
P4 K5 S	97.7
P4 K6 S	97.7
P5 K4 S	97.7
P5 K5 S	98.1
P5 K6 S	97.0
P6 K4 S	98.3
P6 K5 S	98.1
P6 K6 S	98.2
MEAN	97.5

PLOT AREA HARVESTED 0.00210

81/R/RN/17 BARLEY SERIES II

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	5.86
P6 K6 T	7.30
- - S	6.44
P2 - S	6.37
P3 - S	6.39
P4 - S	7.05
P5 - S	6.74
P6 - S	6.95
- K2 S	6.44
- K3 S	6.67
- K4 S	6.37
- K5 S	6.82
- K6 S	6.88
P1 K1 S	5.57
P1 K3 S	6.34
P2 K2 S	6.78
P3 K1 S	6.72
P3 K3 S	6.81
P3 K4 S	6.50
P4 K3 S	7.06
P4 K4 S	6.99
P4 K5 S	7.01
P4 K6 S	7.07
P5 K4 S	6.78
P5 K5 S	7.08
P5 K6 S	6.98
P6 K4 S	7.33
P6 K5 S	7.00
P6 K6 S	6.92
MEAN	6.70

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.374 MIN REP
	0.305 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	11	0.374	5.6

GRAIN MEAN DM% 86.7

PLOT AREA HARVESTED 0.00286

81/R/RN/17

S. BEANS SERIES III

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	3.58
P6 K6 T	3.87
- - S	3.54
P2 - S	3.70
P3 - S	3.60
P4 - S	3.68
P5 - S	3.94
P6 - S	4.82
- K2 S	3.80
- K3 S	3.80
- K4 S	3.60
- K5 S	3.73
- K6 S	4.86
P1 K1 S	2.78
P1 K3 S	3.96
P2 K2 S	2.93
P3 K1 S	3.93
P3 K3 S	3.55
P3 K4 S	3.75
P4 K3 S	3.46
P4 K4 S	3.69
P4 K5 S	4.28
P4 K6 S	3.97
P5 K4 S	4.57
P5 K5 S	4.22
P5 K6 S	4.05
P6 K4 S	4.28
P6 K5 S	4.67
P6 K6 S	4.56
MEAN	3.87

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.604 MIN REP
	0.493 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	11	0.604	15.6

GRAIN MEAN DM% 87.1

PLOT AREA HARVESTED 0.00341

81/R/RN/17

WINTER WHEAT SERIES IV

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	7.67
P6 K6 T	7.11
- - S	7.28
P2 - S	7.56
P3 - S	7.74
P4 - S	7.45
P5 - S	7.90
P6 - S	8.27
- K2 S	7.60
- K3 S	7.19
- K4 S	6.88
- K5 S	7.46
- K6 S	8.13
P1 K1 S	7.18
P1 K3 S	3.43
P2 K2 S	7.54
P3 K1 S	7.87
P3 K3 S	7.06
P3 K4 S	8.36
P4 K3 S	6.88
P4 K4 S	7.30
P4 K5 S	7.41
P4 K6 S	7.71
P5 K4 S	7.85
P5 K5 S	7.06
P5 K6 S	8.14
P6 K4 S	7.66
P6 K5 S	5.89
P6 K6 S	7.77
MEAN	7.38

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.485 MIN REP
	0.396 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	11	0.485	6.6

GRAIN MEAN DM% 78.8

PLOT AREA HARVESTED 0.00286