

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/WW/9 Erynia and Aphid Control - Winter Wheat

Rothamsted Research

Rothamsted Research (1983) *82/R/WW/9 Erynia and Aphid Control - Winter Wheat ; Yields Of The Field Experiments 1982*, pp 241 - 242 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/WW/9

WINTER WHEAT

ERYNIA AND APHID CONTROL

Object: To compare the effects of introducing *Erynia neoaphidis* with two forms and times of applying pirimicarb on cereal aphid population and grain yield - Geescroft.

Sponsors: N. Wilding, G.J.W. Dean.

Design: 3 randomised blocks of 7 plots.

Whole plot dimensions: 6.0 x 6.0.

Treatments:

APH CONT	Chemical and biological aphid control:
NONE	None
EN INTAC	<i>Erynia neoaphidis</i> applied as intact, dried, fungus-killed aphids at 2.5 kg on 15 June and 22 June
EN GRND	<i>Erynia neoaphidis</i> applied as ground, dried, fungus-killed aphids at 2.5 kg on 15 June and 22 June
PS 25 MY	Pirimicarb at 0.14 kg, standard formulation on 25 May
PS 8 JN	Pirimicarb at 0.14 kg, standard formulation on 8 June
PM 25 MY	Pirimicarb at 0.14 kg, microencapsulated on 25 May
PM 8 JN	Pirimicarb at 0.14 kg, microencapsulated on 8 June

Basal applications: Manures: 'Nitro-Chalk' applied at 480 kg. Weedkillers: Mecoprop (as 'Methoxone M' at 5.0 l) with isoproturon at 2.0 kg in 250 l.

Seed: Maris Huntsman, sown at 190 kg.

Cultivations, etc.: - Spring-tine cultivated: 8 Oct, 1981. Heavy spring-tine cultivated twice: 12 Oct. Spring-tine cultivated, rotary harrowed, seed sown: 13 Oct. Weedkillers applied: 15 Apr, 1982. N applied: 23 Apr. Combine harvested: 11 Aug. Previous crops: W. beans 1980, potatoes 1981.

NOTE: Aphid numbers were counted weekly during June and July. Samples of living aphids were taken to determine proportions of those infected with *Erynia*.

82/R/WW/9

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

APH CONT	
NONE	7.09
EN INTAC	7.79
EN GRND	7.92
PS 25 MY	7.54
PS 8 JN	7.82
PM 25 MY	7.49
PM 8 JN	7.07
MEAN	7.53

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	APH CONT
-----	-----
SED	0.652

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.798	10.6
GRAIN MEAN DM%	86.0		
PLOT AREA HARVESTED	0.00139		