

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/WS/1 Insecticides and Alarm Pheromone - Spring Wheat

Rothamsted Research

Rothamsted Research (1983) *82/R/WS/1 Insecticides and Alarm Pheromone - Spring Wheat* ; Yields Of The Field Experiments 1982, pp 245 - 246 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/WS/1

SPRING WHEAT

INSECTICIDES AND ALARM PHEROMONE

Object: To study the effects of an aphid alarm pheromone on the effectiveness of a contact aphicide - Long Hoos III 8.

Sponsors: D.C. Griffiths, J.A. Pickett.

Design: 4 randomised blocks of 6 plots.

Whole plot dimensions: 2.41 x 4.57.

Treatments:

TREATMENT	Aphicide, pheromone treatments and sprayers:
NONE	None
PHEROMON	Aphid alarm pheromone
PERMET C	Permethrin at 0.10 kg, applied by conventional sprayer
PH+PER C	Aphid alarm pheromone plus permethrin applied by conventional sprayer at 0.10 kg
PIRIM C	Pirimicarb at 0.15 kg applied by conventional sprayer
PIRIM E	Pirimicarb at 0.15 kg applied by electrostatic sprayer

NOTE: Treatments were applied on 30 June; insecticides in 60 l of water by electrostatic sprayer, in 460 l of water by conventional sprayer; pheromone at 5 g, in a solvent, by electrostatic sprayer.

Basal applications: Manures: 'Nitro-Chalk' at 450 kg. Weedkillers: Dicamba with mecoprop and MCPA, (as 'Banlene Plus' at 4.9 l) applied with the tridemorph in 340 l. Fungicides: Tridemorph at 0.53 kg, prochloraz at 0.39 kg in 340 l.

Seed: Timmo, sown at 180 kg.

Cultivations, etc.: Ploughed: 29 Jan, 1982. Zig-zag harrowed, power harrowed, seed sown and N applied: 13 Apr. Weedkillers with tridemorph applied: 18 May. Prochloraz applied: 30 June. Combine harvested: 2 Sept. Previous crops: S. barley 1980, s. beans 1981.

NOTE: Aphids were counted five days after spray applications. Plants were analysed for insecticide residues after spraying.

82/R/WS/1

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	NONE	PHEROMON	PERMET C	PH+PER C	PIRIM C	PIRIM E	MEAN
	5.22	4.94	5.16	5.27	4.94	5.04	5.10

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT
SED	0.135

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	15	0.191	3.8

GRAIN MEAN DM% 81.9

PLOT AREA HARVESTED 0.00075