

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/WW/4 Seed Rates and Divided N Dressings - W. Wheat

Rothamsted Research

Rothamsted Research (1984) *83/R/WW/4 Seed Rates and Divided N Dressings - W. Wheat* ; Yields Of The Field Experiments 1983, pp 229 - 231 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/WW/4

WINTER WHEAT

SEED RATES AND DIVIDED N DRESSINGS

Object: To study the effects of a range of rates of early nitrogen dressings on the growth and yield of wheat sown at one third or at standard seed rate - Little Hoos.

Sponsors: J. McEwen, R. Moffitt.

Design: 2 randomised blocks of 30 plots.

Whole plot dimensions: 3.0 x 10.0.

Treatments: All combinations of:-

1. SD RATE Seed rates (kg):

67
200

2. EARLY N Nitrogen fertilizer applied 22 Feb, 1983 (kg N) as 'Nitro-Chalk':

0
25
50
75

3. APRIL N Nitrogen fertilizer applied 13 Apr (kg N) as 'Nitro-Chalk':

75
100
125

plus extra treatments, all combinations of:-

1. SD RATEX Seed rates (kg):

67
200

2. APRIL NX Nitrogen fertilizer applied 13 Apr (kg N):

150
175
200

Basal applications: Manures: (0:18:36) at 420 kg. Weedkillers: Paraquat at 0.83 kg ion in 250 l. Isoproturon at 2.1 kg with dicamba, mecoprop, and MCPA (as 'Poly-Farmon CMPP' at 4.2 l) in 250 l. Fungicides: Carbendazim at 0.15 kg, maneb at 1.6 kg and tridemorph at 0.38 kg in 500 l. Growth regulator: Chlormequat at 1.1 kg in 250 l.

Seed: Avalon.

83/R/WW/4

Cultivations, etc.:— Discd: 25 Aug and 1 Sept, 1982. PK applied: 10 Sept.
Paraquat applied: 15 Oct. Rotary harrowed, seed sown: 27 Oct.
Isoproturon with 'Poly-Farmon CMPP' applied: 11 Mar, 1983. Growth
regulator applied: 29 Apr. Fungicides applied: 13 June. Combine
harvested: 10 Aug. Previous crops: W. wheat 1981, w. oats 1982.

NOTES: (1) Plant counts were made in February, shoot counts in March and
April and ear counts in July.

(2) 1000 grain weights and N content of grain were measured.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

EARLY N	0	25	50	75	MEAN
SD RATE					
67	7.81	8.56	8.64	9.09	8.52
200	7.88	8.62	8.98	9.19	8.67
MEAN	7.85	8.59	8.81	9.14	8.60
APRIL N	75	100	125	MEAN	
SD RATE					
67	8.30	8.60	8.67	8.52	
200	8.16	8.72	9.13	8.67	
MEAN	8.23	8.66	8.90	8.60	
APRIL N	75	100	125	MEAN	
EARLY N					
0	7.19	8.06	8.28	7.85	
25	8.13	8.73	8.90	8.59	
50	8.49	8.81	9.14	8.81	
75	9.10	9.04	9.28	9.14	
MEAN	8.23	8.66	8.90	8.60	
SD RATE	APRIL N	75	100	125	
67	EARLY N				
	0	7.26	8.04	8.12	
	25	8.23	8.80	8.65	
	50	8.65	8.64	8.64	
	75	9.07	8.92	9.27	
200	0	7.13	8.07	8.45	
	25	8.04	8.67	9.15	
	50	8.34	8.97	9.64	
	75	9.12	9.17	9.29	
APRIL NX	150	175	200	MEAN	
SD RATEX					
67	9.00	8.65	9.16	8.94	
200	8.38	9.58	9.23	9.06	
MEAN	8.69	9.12	9.19	9.00	
GRAND MEAN	8.68				

83/R/WW/4

GRAIN TONNES/HECTARE

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	SD RATE	EARLY N	APRIL N	SD RATE EARLY N
SED	0.099	0.139	0.121	0.197

TABLE	SD RATE APRIL N	EARLY N APRIL N	SD RATE EARLY N APRIL N	SDRATES
-------	--------------------	--------------------	-------------------------------	---------

SED	0.171	0.241	0.341	0.197
-----	-------	-------	-------	-------

TABLE	APRIL NX	SD RATES APRIL NX
-------	----------	----------------------

SED	0.241	0.341
-----	-------	-------

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	29	0.341	3.9

GRAIN MEAN DM% 85.8

PLOT AREA HARVESTED 0.00229