

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/BE/8 Control of Stem Nematode - W. Beans

Rothamsted Research

Rothamsted Research (1984) *83/R/BE/8 Control of Stem Nematode - W. Beans ; Yields Of The Field Experiments 1983*, pp 284 - 285 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/BE/8

WINTER BEANS

CONTROL OF STEM NEMATODE

Object: To study the effects of a range of chemicals, rates and times of application on the control of stem nematode (*Ditylenchus dipsaci*) and on the yield of w. beans - Highfield 0 and E III.

Sponsor: A.G. Whitehead.

Design: 3 randomised blocks of 12 plots.

Whole plot dimensions: 2.29 x 4.57.

Treatments:

NEMACIDE	Nematicides, rates and times of application:	
	Applied in the furrow at sowing	Top-dressed after emergence
NONE	None	None (duplicated)
AL1+AL1	Aldicarb at 1.17 kg	+ aldicarb at 1.25 kg
AL2+AL2	Aldicarb at 2.34 kg	+ aldicarb at 2.5 kg
AL4+AL4	Aldicarb at 4.68 kg	+ aldicarb at 5.0 kg
AL2+TH2	Aldicarb at 2.34 kg	+ thiabendazole at 2.5 kg
AL2+TH4	Aldicarb at 2.34 kg	+ thiabendazole at 5.0 kg
CA2+TH2	Carbofuran at 2.38 kg	+ thiabendazole at 2.5 kg
CA2+TH4	Carbofuran at 2.38 kg	+ thiabendazole at 5.0 kg
CA1+CA1	Carbofuran at 1.19 kg	+ carbofuran at 1.25 kg
CA2+CA2	Carbofuran at 2.38 kg	+ carbofuran at 2.5 kg
CA4+CA4	Carbofuran at 4.76 kg	+ carbofuran at 5.0 kg

- NOTES: (1) To ensure the presence of stem nematode, infested straw was spread on the site and ploughed in.
 (2) Thiabendazole was applied in 7,650 l water.
 (3) Post emergence treatments were applied to w. beans on 25 May, 1983

Basal applications: Manures: (0:20:20) at 630 kg. Weedkillers: Simazine at 1.1 l in 560 l. Fungicide: Benomyl at 0.56 kg in 560 l, on two occasions, with the insecticide on the second. Insecticide: Pirimicarb at 0.14 kg on three occasions, with the fungicide on the first, in 560 l on the second and third.

Seed: Throws MS, sown at 240 kg.

Cultivations, etc.: PK applied, infested straw applied, ploughed in: 17 Aug, 1982. Rotary harrowed, seed sown: 11 Nov. Weedkiller and fungicide applied: 4 Feb, 1983. Fungicide with insecticide applied: 10 June. Insecticide applied: 30 June, 19 July. Harvested by hand: 3 Aug. Previous crops: S. beans 1981, w. beans 1982.

NOTE: The percentage of stems infested with stem nematode was assessed in July.

83/R/BE/8

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

NEMACIDE	
NONE	0.81
AL1+AL1	1.30
AL2+AL2	1.78
AL4+AL4	2.21
AL2+TH2	1.45
AL2+TH4	1.71
CA2+TH2	1.86
CA2+TH4	1.67
CA1+CA1	2.14
CA2+CA2	2.44
CA4+CA4	2.80
MEAN	1.75

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	NEMACIDE	

SED	0.367	MIN REP
	0.318	MAX-MIN

NEMACIDE
 MAX-MIN NONE V ANY OF THE REMAINDER
 MIN REP ANY OF THE REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	23	0.449	25.7
GRAIN MEAN DM%	78.6		
PLOT AREA HARVESTED	0.00035		