

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



Winter Oilseed Rape

Rothamsted Research

Rothamsted Research (1984) *Winter Oilseed Rape* ; Yields Of The Field Experiments 1983, pp 310 - 313 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/RA/1

WINTER OILSEED RAPE

MUSTARD OILS AND LIGHT LEAF SPOT

Object: To study the effects of mustard oils, sulphur and benomyl on the incidence of light leaf spot (*Pyrenopeziza brassicae*) and on the yield of two w. oilseed rape varieties with contrasting glucosinolate levels - Long Hoos V 4.

Sponsors: C.J. Rawlinson, G.R. Cayley, J.A. Pickett.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 1.83 x 6.10.

Treatments: All combinations of:

1. VARIETY Varieties:

DP 1305
NORLI

2. FUNGICIDE Fungicides:

NONE	None
BENOMYL	Benomyl at 0.5 kg in 4.2 l
M OIL 1	Mustard oil 1 at 0.15 kg in 4.2 l
M OIL 2	Mustard oil 2 at 0.24 kg in 4.2 l
SULPHUR	Elemental sulphur at 10 kg in 340 l (as 'Thiovit')

NOTES: (1) All fungicide treatments were applied on 30 Nov, 1982 and 12 Apr, 1983; sulphur was applied by hydraulic sprayer, the remainder by electrostatic sprayer.

(2) Diseases were so severe on VARIETY DP1305 that all plants, irrespective of treatments were killed by the end of April.

Basal applications: Manures: 'Nitro-Chalk' at 900 kg. Weedkillers: Benazolin ethyl ester at 0.67 kg with 3, 6-dichloropicolinic acid at 0.11 kg in 450 l. Desiccant: Diquat at 0.84 kg ion in 250 l.

Seed: Sown at 19 kg.

Cultivations, etc.: Spring-tine cultivated twice: 25 Aug, 1982. Seed sown by hand: 26 Aug. Weedkillers applied: 15 Oct. N applied: 14 Mar, 1983. Desiccant applied: 25 July. Combine harvested: 27 July. Previous crops: Potatoes 1981, fallow 1982.

NOTE: Phoma and light leaf spot were assessed on five occasions during the season.

83/R/RA/1

NORLI ONLY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGCIDE	NONE	BENOMYL	M OIL 1	M OIL 2	SULPHUR	MEAN
	2.05	2.60	2.29	2.47	2.46	2.37

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGCIDE
-----	-----
SED	0.246

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	8	0.301	12.7

GRAIN MEAN DM% 75.4

PLOT AREA HARVESTED 0.00111

83/R/RA/4

WINTER OILSEED RAPE

CONTROL OF PESTS AND DISEASES

Object: To study the effects of carbofuran, cypermethrin, benomyl and triazophos on the incidence of pests and diseases and on the yield of w. oilseed rape - Bones Close.

Sponsors: R.J. Tuppen (MAFF), G.J.W. Dean, I.H. Williams, C.J. Rawlinson.

Design: 4 randomised blocks of 6 plots.

Whole plot dimensions: 4.0 x 20.0.

Treatments: All combinations of:-

- | | |
|-------------|--|
| 1. INSCTCDE | Insecticides: |
| NONE | None |
| CARBOFUR | Carbofuran at 2.5 kg as granules on 3 Sept, 1982 |
| CYPERMET | Cypermethrin at 0.035 l in 500 l on 3 June, 1983 |
| 2. FUNG INS | Fungicide and additional insecticide: |
| NONE | None |
| BEN+TRI | Benomyl at 0.50 kg on 15 Nov, 1982, and triazophos on 8 June, 1983 |

NOTE: 'Actipron', a wetting agent, was applied at 2.5 l with the benomyl.

Basal applications: Manures: (5:14:30) at 450 kg. 'Nitro-Chalk' at 500 kg followed by 450 kg. Weedkillers: Paraquat at 0.70 kg ion in 250 l. Diquat at 0.60 kg ion applied with 'Agral', a wetting agent, at 0.5 l in 500 l. Propyzamide and 3,6 - dichloropicolinic acid (as 'Matri-Kerb' at 1.63 kg) in 250 l.

Seed: Jet Neuf, dressed thiram and thiabendazole, sown at 7.8 kg.

Cultivations, etc.: Paraquat applied: 23 Aug, 1982. Discd: 29 Aug. NPK applied, seed sown: 31 Aug. 'Matri-Kerb' applied: 9 Nov. First N applied: 23 Feb, 1983. Second N applied: 28 Mar. Diquat applied: 22 July. Combine harvested: 28 July. Previous crops: W. wheat 1981, w. wheat and w. barley 1982.

NOTE: Light leaf spot was assessed in early spring, May and June.

83/R/RA/4

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNG INS INSCDCE	NONE	BEN+TRI	MEAN
NONE	3.17	3.39	3.28
CARBOFUR	3.11	3.53	3.32
CYPERMET	2.91	3.51	3.21
MEAN	3.06	3.47	3.27

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	INSCDCE	FUNG INS	INSCDCE FUNG INS
SED	0.219	0.179	0.310

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	15	0.439	13.4

GRAIN MEAN DM% 85.6

PLOT AREA HARVESTED 0.00640