

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/P/1 Varieties and Stem Canker - Potatoes

Rothamsted Research

Rothamsted Research (1984) *83/R/P/1 Varieties and Stem Canker - Potatoes* ; Yields Of The Field Experiments 1983, pp 314 - 317 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/P/1

POTATOES

VARIETIES AND STEM CANKER

Object: To study the effects of stem canker (*Rhizoctonia solani*) on plant development and yield of a range of early and maincrop potato varieties - Summerdells II.

Sponsors: G.A. Hide, P.J. Read.

Design: Early varieties: 3 randomised blocks of 20 plots.
Maincrop varieties: 3 randomised blocks of 28 plots.

Whole plot dimensions: 3.00 x 9.14.

Treatments:

To EARLY varieties, all combinations of:-

- | | |
|-------------|---|
| 1. VARIETY | Varieties: |
| A COMET | Arran Comet |
| ESTIMA | Estima |
| M PEER | Maris Peer |
| U PRINCE | Ulster Prince |
| U SCEPTR | Ulster Sceptre |
| 2. RHZ FUNG | R. solani infection and fungicide to tubers: |
| NONE | None |
| SEED R | Seed with black scurf symptom |
| SEED R+F | Seed with black scurf symptom, treated with iprodione |
| SOIL R | R. solani inoculum to soil |

To MAINCROP varieties, all combinations of:-

- | | |
|-------------|---|
| 1. VARIETY | Varieties: |
| CARA | Cara |
| DESIREE | Desiree |
| K EDWARD | King Edward |
| P CROWN | Pentland Crown |
| M PIPER | Maris Piper |
| P SQUIRE | Pentland Squire |
| RECORD | Record |
| 2. RHZ FUNG | R. solani infection & fungicide to tubers: |
| NONE | None |
| SEED R | Seed with black scurf symptom |
| SEED R+F | Seed with black scurf symptom, treated with iprodione |
| SOIL R | R. solani inoculum to soil |

NOTES: (1) The fungal inoculum was grown on a vermiculite plus malt medium. It was applied by hand around the tubers at planting.
(2) The iprodione treatment was applied by dipping tubers into a 0.05% solution for five minutes.

83/R/P/1

Basal applications: Manures: (0:18:36) at 700 kg. (10:10:15+4.5 Mg) at 2010 kg. Weedkillers: Paraquat at 0.84 kg ion in 250 l. Linuron at 1.0 kg with paraquat at 0.56 kg ion in 500 l. Fungicides: Mancozeb at 1.4 kg in 250 l on three occasions. Fentin hydroxide at 0.28 kg in 250 l applied on five occasions, with the insecticide on the first four occasions. Insecticide: Pirimicarb at 0.14 kg. Desiccant: BOV at 170 l.

Cultivations, etc:- PK applied: 20 Sept, 1982. Paraquat applied: 16 Oct. Ploughed: 9 Dec. NPK Mg applied: 4 May, 1983. Rotary harrowed for early-sown potatoes, heavy spring-tine cultivated for maincrop: 11 May. Early potatoes planted by hand, maincrop area rotary harrowed: 16 May. Maincrop potatoes planted by hand: 23 May. Linuron and paraquat applied: 7 June. Mancozeb applied: 22 June. Fentin hydroxide with pirimicarb applied: 1 July, 8 July, 18 July, 28 July. Fentin hydroxide alone applied: 11 Aug. Mancozeb applied: 25 Aug, 9 Sept. Haulm mechanically destroyed: 7 Oct. Haulm desiccant applied: 19 Oct. Lifted: 26 Oct. Previous crops: S. barley 1981, w. oats 1982.

- NOTES: (1) Emergence counts were made in June.
(2) Measurements of aspects of plant growth and of *R. solani* infection of stem bases and stolons were made in July and August.
(3) Infection of mature tubers was assessed.

83/R/P/1 EARLY POTATOES

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

RHZ FUNG VARIETY	NONE	SEED R	SEED R+F	SOIL R	MEAN
A COMET	29.9	30.4	29.6	30.5	30.1
ESTIMA	26.2	29.7	27.7	27.9	27.9
M PEER	15.8	15.2	16.8	15.4	15.8
U PRINCE	16.7	16.7	16.4	16.9	16.7
U SCEPT	17.1	17.9	17.8	18.1	17.7
MEAN	21.1	22.0	21.7	21.8	21.6

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY	RHZ FUNG	VARIETY RHZ FUNG
SED	0.76	0.68	1.52

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	38	1.86	8.6

PERCENTAGE WARE 4.44CM (1.75 INCH) RIDDLE

***** TABLES OF MEANS *****

RHZ FUNG VARIETY	NONE	SEED R	SEED R+F	SOIL R	MEAN
A COMET	61.3	65.8	56.0	71.2	63.5
ESTIMA	68.5	71.5	69.6	72.7	70.6
M PEER	31.0	37.9	32.4	40.2	35.4
U PRINCE	74.9	81.1	81.0	81.1	79.5
U SCEPT	54.5	55.2	53.1	61.2	56.0
MEAN	58.0	62.3	58.4	65.3	61.0

PLOT AREA HARVESTED 0.00117

83/R/P/1 MAIN CROP POTATOES

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

RHZ FUNG VARIETY	NONE	SEED R	SEED R+F	SOIL R	MEAN
CARA	34.1	37.0	37.0	37.3	36.3
DESIREE	32.6	32.6	33.7	31.2	32.5
K EDWARD	36.3	36.6	34.9	31.8	34.9
P CROWN	33.4	33.7	31.4	34.6	33.2
M PIPER	32.0	33.0	31.4	31.6	32.0
P SQUIRE	38.3	36.4	36.8	36.3	36.9
RECORD	29.5	26.4	27.4	26.7	27.5
MEAN	33.7	33.7	33.2	32.8	33.4

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY	RHZ FUNG	VARIETY RHZ FUNG
SED	0.89	0.67	1.77

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	54	2.17	6.5

PERCENTAGE WARE 4.44CM (1.75 INCH) RIDDLE

***** TABLES OF MEANS *****

RHZ FUNG VARIETY	NONE	SEED R	SEED R+F	SOIL R	MEAN
CARA	84.9	88.3	85.8	86.3	86.3
DESIREE	62.8	69.4	63.3	69.8	66.3
K EDWARD	42.7	46.2	35.9	55.1	45.0
P CROWN	80.2	84.6	79.9	85.6	82.6
M PIPER	49.8	51.3	51.0	54.6	51.6
P SQUIRE	81.7	85.9	84.0	88.9	85.1
RECORD	56.4	59.3	63.4	60.9	60.0
MEAN	65.5	69.3	66.2	71.6	68.1

PLOT AREA HARVESTED 0.00117