

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/CS/10 and 83/W/CS/10 Long-term Liming - Potatoes

Rothamsted Research

Rothamsted Research (1984) *83/R/CS/10 and 83/W/CS/10 Long-term Liming - Potatoes* ; Yields Of The Field Experiments 1983, pp 96 - 99 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/CS/10 and 83/W/CS/10

LONG TERM LIMING

Object: To study the effects of different amounts of lime on the yields of a sequence of crops. The effects of P are also studied - Rothamsted (R) Sawyers I and Woburn (W) Stackyard C.

Sponsors: S.P. McGrath, D.P. Stribley.

The 22nd year, potatoes.

For previous years see 'Details' 1967, 1973 and 74-82/R&W/CS/10.

Design: 2 randomised blocks of 16 plots.

Whole plot dimensions: 6.40 x 18.3.

Treatments: All combinations of:-

1. CHALK Ground chalk (tonnes CaCO₃) (total applied 1962-83):

R	W	Rothamsted Total			Woburn Total		
		1962-78	1982	1983	1962-78	1982	1983
0	0	0	0	0	0	0	0
14	8	7	2	5	6	2	0
23	24	15	5	3	14	5	5
50	43	30	10	10	23	10	10

2. P P fertilizer applied:

	Until 1978	1981	1982	1983
	R & W	R & W	R & W	R & W
0	0	0	0	0
P1	0	P1	P1	0
P2	P	P1	0	P2
P3	P	P3	P1	P2

Rates 1981-1983 P1, P2, P3, P4 = 25, 50, 75, 100 kg P as superphosphate

- NOTES: (1) Until 1978 test P was applied cumulatively, rates varied with crop. None was applied in 1979 and 1980 (fallow). K was applied, cumulatively, to P1 and P3 plots. Since 1981 K has been applied basally.
- (2) A sub plot test of Mg applied in earlier years has been ignored. Since 1981 Mg has been applied basally.

Basal applications:

Sawyers I (R): Manures: N at 250 kg as 'Nitro-Chalk', K at 210 kg as muriate of potash, Mg at 40 kg as kieserite. Weedkillers: Linuron at 1.0 l with paraquat at 0.4 kg ion in 500 l. Fungicide: Mancozeb at 1.4 kg in 250 l on three occasions. Fentin hydroxide at 0.28 kg in 250 l on five occasions with the insecticide on the first four occasions. Insecticide: Pirimicarb at 0.14 kg. Haulm desiccant: Undiluted BOV at 170 l.

83/R/CS/10 and 83/W/CS/10

Stackyard C (W): Manures: N at 260 kg as 'Nitro-Chalk', K at 210 kg as muriate of potash, Mg at 40 kg as kieserite. Weedkillers: Glyphosate at 1.5 kg in 280 l. Linuron at 1.0 l with paraquat at 0.4 kg ion in 250 l. Fungicide: Mancozeb at 1.4 kg in 250 l on four occasions with the insecticide on the first, third and fourth occasions. Fentin hydroxide at 0.28 kg in 250 l on four occasions with insecticide on the first three occasions. Insecticide: Pirimicarb at 0.14 kg. Haulm desiccant: Diquat at 0.50 kg ion in 250 l.

Seed: Pentland crown.

Cultivations, etc.:-

Sawyers I (R): Ground chalk treatments applied: 26 Nov, 1982. Ploughed: 1 Dec. P treatments and N, K and Mg applied: 11 May, 1983. Rotary harrowed, potatoes planted: 23 May. Rotary ridged: 26 May. Weedkillers applied: 7 June. Mancozeb applied: 22 June, 25 Aug, 9 Sept. Fentin hydroxide applied with insecticide: 1 July, 8 July, 18 July, 28 July. Fentin hydroxide applied: 11 Aug. Haulm mechanically destroyed: 11 Oct. Haulm desiccant applied: 19 Oct. Lifted: 28 Oct.

Stackyard C (W): Glyphosate applied: 12 Sept, 1982. Ploughed: 8 Oct. Ground chalk treatments applied: 4 Nov. Ploughed: 23 Dec. Heavy spring-tine cultivated: 21 Mar, 1983. P treatments and N, K and Mg applied: 22 Mar. Heavy spring-tine cultivated: 28 Mar, 11 May. Rotary cultivated, potatoes planted: 11 May. Rotary ridged, weedkiller applied: 7 June. Mancozeb applied with insecticide: 22 June, 8 July, 18 July. Mancozeb applied: 1 July. Fentin hydroxide applied with insecticide: 29 July, 11 Aug, 26 Aug. Fentin hydroxide applied: 9 Sept. Haulm mechanically destroyed: 5 Oct. Haulm desiccant applied: 17 Oct. Lifted: 7 Nov.

NOTE: Soil samples were taken from both sites after harvest for chemical analyses.

83/R/CS/10 SAWYERS (R)

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

P	0	P1	P2	P3	MEAN
CHALK					
0	19.3	18.7	27.9	29.4	23.8
14	26.8	28.8	29.4	33.1	29.5
23	26.6	29.2	30.7	33.5	30.0
50	26.3	27.8	29.0	32.2	28.8
MEAN	24.8	26.1	29.2	32.0	28.0

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHALK	P	CHALK P
SED	0.99	0.99	1.97

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	15	1.97	7.0

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

P	0	P1	P2	P3	MEAN
CHALK					
0	97.4	96.6	97.3	97.5	97.2
14	97.6	98.2	96.7	97.3	97.5
23	97.6	97.8	97.6	97.8	97.7
50	97.6	97.5	97.2	98.2	97.6
MEAN	97.5	97.5	97.2	97.7	97.5

PLOT AREA HARVESTED 0.00274

83/W/CS/10 STACKYARD (W)

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

P	0	P1	P2	P3	MEAN
CHALK					
0	29.8	40.2	45.5	43.0	39.6
8	43.9	52.2	46.1	50.0	48.1
24	37.7	42.9	41.9	42.4	41.2
43	36.6	41.9	37.5	40.0	39.0
MEAN	37.0	44.3	42.7	43.8	42.0

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHALK	P	CHALK P
SED	0.88	0.88	1.77

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	15	1.77	4.2

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

P	0	P1	P2	P3	MEAN
CHALK					
0	97.6	96.8	95.7	95.9	96.5
8	97.8	96.9	96.9	96.2	97.0
24	97.5	98.2	96.0	96.1	97.0
43	98.6	98.5	97.4	97.6	98.0
MEAN	97.9	97.6	96.5	96.5	97.1

PLOT AREA HARVESTED 0.00274