

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/WW/7 Aphid Control by Erynia - W. Wheat

Rothamsted Research

Rothamsted Research (1985) *84/R/WW/7 Aphid Control by Erynia - W. Wheat ; Yields Of The Field Experiments 1984*, pp 234 - 235 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/WW/7

WINTER WHEAT

APHID CONTROL BY ERYNIA

Object: To study the effect on caged and uncaged aphid populations of applying two amounts of the aphid pathogenic fungus *Erynia neoaphidis* on the incidence of aphids and on the yield of w. wheat - Long Hoos V 4.

Sponsor: N. Wilding.

Design: 3 randomised blocks of 6 plots.

Whole plot dimensions: 2.14 x 2.0.

Treatments: All combinations of:-

- | | |
|-------------|---|
| 1. COVER | Covering on plots: |
| NONE | None |
| CAGED | Mesh-sided cages covering plots from 18 May, 1984 to 29 June. |
| 2. INOCULUM | Rate of inoculum: |
| NONE | None |
| E NEO 1 | E. neoaphidis applied as a powder of mummified aphids at 0.5 kg on 17 June and 4 July |
| E NEO 2 | E. neoaphidis applied as a powder of mummified aphids at 5.0 kg on 17 June and 4 July |

Note: *Sitobion avenae* and *Metopolophium dirhodum* were released on all plots, during a ten-day period from the end of May.

Basal applications: Manures: Chalk at 2.9 t. 'Nitro-Chalk' at 450 kg. Weedkillers: Glyphosate at 1.4 kg in 250 l. Mecoprop, bromoxynil and ioxynil (as 'Brittox' at 3.5 l) in 220 l applied with the fungicide. Fungicide: Triadimefon at 0.12 kg in 220 l on two occasions, the first with the 'Brittox'. Insecticide: Pyrethrum dust (as 'Anti-ant' at 116 kg).

Seed: Maris Huntsman, sown at 190 kg.

Cultivations, etc.: Glyphosate applied: 18 Aug, 1983. Chalk applied: 23 Aug. Ploughed: 31 Aug. Spring-tine cultivated, seed sown: 5 Oct. 'Brittox' with the fungicide applied, N applied: 10 Apr, 1984. Fungicide alone applied: 4 May. Pyrethrum applied: 21 May. Combine harvested: 22 Aug. Previous crops: Fallow 1982, w. rape 1983.

NOTE: Samples of live aphids were taken weekly from mid June until late July to determine proportions infected with *Erynia*, and the total population was estimated weekly.

84/R/WW/7

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

INOCULUM COVER	NONE	E NEO 1	E NEO 2	MEAN
NONE	7.00	6.88	7.07	6.98
CAGED	4.04	3.61	3.79	3.82
MEAN	5.52	5.24	5.43	5.40

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	INOCULUM	COVER	INOCULUM COVER
SED	0.232	0.190	0.328

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	10	0.402	7.5

GRAIN MEAN DM% 86.9

PLOT AREA HARVESTED 0.00010