

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/RA/6 Spring Nitrogen Inhibitors - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1985) *84/R/RA/6 Spring Nitrogen Inhibitors - W. Oilseed Rape* ; Yields Of The Field Experiments 1984, pp 316 - 317 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/RA/6

WINTER OILSEED RAPE

SPRING NITROGEN INHIBITORS

Object: To study the effects of adding nitrification inhibitors to prilled urea, applied in spring, on the yield and nitrogen uptake of w. oilseed rape - Whittlocks.

Sponsors: G.A. Rodgers, A. Penny, M.V.Hewitt.

Design: 2 randomised blocks of 9 plots.

Whole plot dimensions: 4.0 x 20.0.

Treatments: All combinations of:-

- | | |
|------------|---|
| 1. N INHIB | Forms of nitrogen and nitrification inhibitors: |
| AN O | Ammonium nitrate (as 'Nitro-Chalk'), no inhibitor |
| PU O | Prilled urea, no inhibitor |
| PU DIC | Prilled urea and dicyandiamide at 33 kg |
| PU HYD | Prilled urea and hydroquinone at 13 kg |

- | | |
|-------------|-------------------------------------|
| 2. SPRING N | Nitrogen rates (kg N) and times: |
| 75E+75L | 75 on 10 Feb and 75 on 21 Mar, 1984 |
| 150M | 150 on 5 Mar |

plus one extra treatment

EXTRA

NONE No spring nitrogen fertilizer or inhibitor

Basal applications: Manures: 'Nitro-Chalk' at 220 kg. Weedkillers: TCA at 11 kg in 250 l. Propyzamide with 3, 6-dichloropicolinic acid (as 'Matrikerb' at 1.6 kg) in 500 l. Fungicide: Prochloraz at 0.50 kg in 250 l. Desiccant: Diquat at 0.6 kg ion with 'Agral', a wetting agent, at 0.5 l, in 500 l.

Seed: Jet Neuf, dressed thiram, fenpropimorph and iprodione, sown at 9.0 kg.

Cultivations, etc.: - Discd: 17 Aug, 1983. N applied: 22 Aug. TCA applied: 23 Aug. Spring-tine cultivated: 24 Aug. Seed direct drilled: 25 Aug. 'Matrikerb' applied: 25 Oct. Fungicide applied: 18 Apr, 1984. Desiccant applied: 25 July. Combine harvested: 30 July. Previous crops: W. wheat 1982, w. barley 1983.

- NOTES: 1) Ammonia volatilisation was measured after application of N dressings.
2) Soil samples were taken from February to June for N analyses.
3) Oil and protein contents of grain were measured.

84/R/RA/6

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** TABLES OF MEANS *****

SPRING N	75E+75L	150M	MEAN
N INHIB			
AN O	3.09	3.41	3.25
PU O	3.17	3.11	3.14
PU DIC	2.96	3.16	3.06
PU HYD	3.33	3.29	3.31
MEAN	3.14	3.24	3.19

NONE 1.77

GRAND MEAN 3.03

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	N INHIB	SPRING N	N INHIB SPRING N & NONE
SED	0.109	0.077	0.155

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	8	0.155	5.1

GRAIN MEAN DM% 88.9

PLOT AREA HARVESTED 0.00465