

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1984

[Full Table of Content](#)



### 84/R/RA/8 Mustard Oil - W. Oilseed Rape

#### Rothamsted Research

Rothamsted Research (1985) *84/R/RA/8 Mustard Oil - W. Oilseed Rape ; Yields Of The Field Experiments 1984*, pp 320 - 321 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/RA/8

WINTER OILSEED RAPE

MUSTARD OIL

Object: To study the effects of carbendazim, prochloraz and a synthetic mustard oil formulation on fungal diseases and yield of w. oilseed rape - Whittlocks.

Sponsors: C.J. Rawlinson, G.R. Cayley, J.A. Pickett.

Design: 8 randomised blocks of 4 plots.

Whole plot dimensions: 2.0 x 6.0.

Treatments:

| FUNGICIDE | Fungicides:                                       |
|-----------|---------------------------------------------------|
| NONE      | None                                              |
| MSTD OIL  | Mustard oil at 0.25 kg                            |
| PROCHLOR  | Prochloraz at 0.40 kg                             |
| PROC+CAR  | Prochloraz at 0.40 kg plus carbendazim at 0.15 kg |

NOTES: (1) Treatment sprays were applied to four blocks on 4 Apr, 1984 to the other four on 13 Apr.  
(2) Treatments were applied by electrostatic sprayer in 9.3 l water.

Basal applications: Manures: 'Nitro-Chalk' at 220 kg on the first occasion, and at 460 kg on the second and third occasions. Weedkillers: TCA at 11 kg in 250 l. Propyzamide and 3, 6-dichloropicolinic acid (as 'Matrikerb' at 1.63 kg) in 500 l. Desiccant: Diquat at 0.60 kg ion with 'Agral', a wetting agent, at 0.5 l, in 500 l.

Seed: Jet Neuf, seed dressed gamma HCH, thiram, fenpropimorph and iprodione, sown at 9 kg.

Cultivations, etc.: Discd: 17 Aug, 1983. N applied: 22 Aug. TCA applied: 23 Aug. Spring-tine cultivated: 24 Aug. Seed direct drilled: 25 Aug. 'Matrikerb' applied: 25 Oct. N applied: 15 Feb, 1984. N applied: 5 Apr. Desiccant applied: 25 July. Combine harvested: 31 July. Previous crops: W. wheat 1982, w. barley 1983.

NOTE: Disease incidence and severity, to all parts of the plant, was assessed fortnightly from April to July.

84/R/RA/8

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

| FUNGCIDE | NONE | MSTD OIL | PROCHLOR | PROC+CAR | MEAN |
|----------|------|----------|----------|----------|------|
|          | 4.56 | 4.63     | 4.59     | 4.65     | 4.61 |

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

| TABLE | FUNGCIDE |
|-------|----------|
| ----- | -----    |
| SED   | 0.161    |

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

| STRATUM  | DF | SE    | CV% |
|----------|----|-------|-----|
| BLOCK.WP | 21 | 0.321 | 7.0 |

GRAIN MEAN DM% 83.7

PLOT AREA HARVESTED 0.00120