

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/R/ON/1 Control of Stem Nematode - Onions

Rothamsted Research

Rothamsted Research (1986) *85/R/ON/1 Control of Stem Nematode - Onions ; Yields Of The Field Experiments 1985*, pp 344 - 345 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/ON/1

ONIONS

CONTROL OF STEM NEMATODE

Object: To study the effects of rates, times and methods of applying nematicides on the control of stem nematode and on the yield of onions - Fosters 0 and E VI.

Sponsor: A.G. Whitehead.

Design: 2 randomised blocks of 15 plots.

Whole plot dimensions: 2.29 x 7.32.

Treatments: All combinations of:-

1. NEM S Nematicides to seed furrow:
 ALDICARB Aldicarb
 CARBOFUR Carbofuran
2. NEM S RT Rates of nematicides to seed furrow (kg):
 1
 2
3. NEM F Foliar nematicide, applied by electrostatic sprayer, at
 1.5 kg:
 NONE None
 THIABEND Thiabendazole
 THIODICA Thiodicarb

plus one extra treatment:

EXTRA

NONE None (triplicated)

NOTE: The foliar nematicides were applied in 5.7 l on 2 June, 1985.

Basal applications: Manures: (13:13:20) at 1880 kg. Weedkillers:
Glyphosate at 1.4 kg in 280 l. Propachlor at 4.4 kg in 280 l.

Seed: Robusta, dressed benomyl and thiram, sown at 6.7 kg.

Cultivations, etc.: - Glyphosate applied: 25 Sept, 1984. Ploughed:
16 Nov. NPK applied: 11 Mar, 1985. Rotary harrowed: 13 Mar. Seed
sown, seed furrow treatments applied: 14 Mar. Propachlor applied:
21 Mar. Lifted: 24 Sept. Previous crops: S. oats 1983, w. oats
1984.

NOTE: Stem nematode damage to onions stored after harvest was assessed.

85/R/ON/1

FRESH WEIGHT TONNES/HECTARE

***** TABLES OF MEANS *****

NEM S RT	1	2	MEAN
NEM S			
ALDICARB	54.1	56.4	55.3
CARBOFUR	44.0	53.6	48.8
MEAN	49.1	55.0	52.0

NEM F	NONE	THIABEND	THIODICA	MEAN
NEM S				
ALDICARB	55.3	55.3	55.2	55.3
CARBOFUR	47.4	49.8	49.2	48.8
MEAN	51.4	52.5	52.2	52.0

NEM F	NONE	THIABEND	THIODICA	MEAN
NEM S RT				
1	44.9	50.8	51.5	49.1
2	57.8	54.3	53.0	55.0
MEAN	51.4	52.5	52.2	52.0

NEM S RT	1	2	3	4	5	6
NEM F	NONE	THIABEND	THIODICA	NONE	THIABEND	THIODICA
NEM S						
ALDICARB	48.9	55.4	58.0	61.7	55.1	52.4
CARBOFUR	40.8	46.1	45.0	53.9	53.5	53.5

EXTRA NONE 0.5

GRAND MEAN 41.7

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	NEM S	NEM S RT	NEM F	NEM S NEM S RT
SED	2.52	2.52	3.09	3.56

TABLE	NEM S NEM F	NEM S RT NEM F	NEM S NEM S RT NEM F
SED	4.37	4.37	6.17

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	11	6.17	11.9
PLOT AREA HARVESTED	0.00084		