

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/R/RN/17 Rates of P and K to the Subsoil - S. Beans, W. Wheat, Potatoes, S. Barley

Rothamsted Research

Rothamsted Research (1986) *85/R/RN/17 Rates of P and K to the Subsoil - S. Beans, W. Wheat, Potatoes, S. Barley* ; Yields Of The Field Experiments 1985, pp 85 - 92 - **DOI:**
<https://doi.org/10.23637/ERADOC-1-19>

85/R/RN/17

RATES OF P AND K TO THE SUBSOIL

Object: To study the effects of a range of rates and frequencies of application of P and K to the subsoil, singly and together, on the yields and nutrient uptakes of a rotation of crops - Meadow.

Sponsors: J. McEwen, A.E. Johnston.

The fifth year, potatoes, s. barley, s. beans, w. wheat.

For previous years see 81-84/R/RN/17.

Design: 4 series (for crops) each of 40 plots.

Whole plot dimensions: 3.0 x 14.0.

Treatments to each series:

TREATMENT Extra P and K and primary cultivation tool in autumn 1980 only, except on A plots, treatments repeated annually, and F plots treatments repeated four yearly:

	P205(kg)	K20(kg)	Tool	
- - -	0	0	Plough	(duplicated)
P6 K6 T	1000	500 to topsoil	"	(")
- - S	0	0	Wye double-digger	(triplicated)
- - SA	0	0	" "	(duplicated)
- - SF	0	0	" "	"
P2 - SA	63	0 to subsoil	" "	"
P3 - SF	125	0 " "	" "	"
P4 - S	250	0 " "	" "	"
P5 - S	500	0 " "	" "	"
P5 - SF	500	0 " "	" "	"
P6 - S	1000	0 " "	" "	"
- K2 SA	0	31 " "	" "	"
- K3 SF	0	63 " "	" "	"
- K4 S	0	125 " "	" "	"
- K5 S	0	250 " "	" "	"
- K5 SF	0	250 " "	" "	"
- K6 S	0	350 " "	" "	"
P1 K1 SA	31	16 " "	" "	"
P1 K3 SA	31	63 " "	" "	"
P2 K2 SA	63	31 " "	" "	"
P3 K1 SA	125	16 " "	" "	"
P3 K3 SA	125	63 " "	" "	"
P3 K4 SF	125	125 " "	" "	"
P4 K3 SF	250	63 " "	" "	"
P4 K4 S	250	125 " "	" "	"
P4 K5 S	250	250 " "	" "	"
P4 K5 SF	250	250 " "	" "	"
P4 K6 S	250	350 " "	" "	"
P5 K4 S	500	125 " "	" "	"
P5 K4 SF	500	125 " "	" "	"
P5 K5 S	500	250 " "	" "	"
P5 K6 S	500	350 " "	" "	"
P6 K4 S	1000	125 " "	" "	"
P6 K5 S	1000	250 " "	" "	"
P6 K6 S	1000	350 " "	" "	"

85/R/RN/17

- NOTES: (1) Subsoiling was done with the Wye double-digger which turns a furrow with a conventional plough share, to a depth of 23 cm, and at the same time rotary cultivates the bottom of the adjacent furrow to a further depth of 15 cm. When applying P and K this was distributed ahead of the rotary cultivator.
- (2) The topsoil PK dressing was equally divided before and after ploughing.
- (3) All plots were conventionally ploughed each autumn unless the Wye double-digging treatment was due.
- (4) The rate of 350 kg K2O applied was in error for 500 kg K2O.

Standard applications:

- Potatoes: Manures: Chalk at 5.0 t. (10:10:15+4.5 Mg) at 1960 kg.
Weedkillers: Linuron at 1.3 kg with paraquat at 0.50 kg ion in 500 l. Fungicides: Mancozeb at 1.4 kg in 200 l on four occasions, with the insecticide on the second and third. Fentin hydroxide at 0.28 kg in 200 l on two occasions. Insecticide: Pirimicarb at 0.14 kg on two occasions.
- S. barley: Manures: Chalk at 5.0 t. (20:10:10) at 630 kg.
Weedkillers: Clopyralid at 0.05 kg with bromoxynil octanoate at 0.24 kg, mecoprop (as 'CMPP' at 3.0 l) applied with the fungicide in 200 l. Fungicide: Tridemorph at 0.52 kg.
- S. beans: Manures: Chalk at 5.0 t. Weedkiller: Simazine at 1.2 kg in 200 l. Fungicide: Benomyl at 0.50 kg, applied with the pirimicarb and a wetting agent ('Agral' at 0.075 l) in 500 l. Insecticides: Phorate at 5.6 kg. Pirimicarb at 0.14 kg.
- W. wheat. Manures: Chalk at 5.0 t. (0:18:36) at 350 kg. 'Nitro-Chalk' (27.5% N) at 540 kg. Weedkillers: Clopyralid at 0.05 kg with bromoxynil octanoate at 0.24 kg and mecoprop (as 'CMPP' at 3.0 l), applied with the tridemorph in 200 l. Fungicides: Tridemorph at 0.52 kg. Propiconazole at 0.25 kg with carbendazim and maneb (as 'Septal' at 2.5 kg) in 200 l. Insecticide: Pirimicarb at 0.14 kg in 200 l.

Seed: Potatoes: Pentland Crown.

S. barley: Klaxon, sown at 160 kg.

S. beans: Minden, sown at 240 kg.

W. wheat: Avalon, sown at 200 kg.

Cultivations, etc.:-

All crops: Chalk applied: 2 Oct, 1984. Treatments applied by double-digger: 14-19 Nov. Ploughed: 26 Nov. Spring-tine cultivated (twice for w. wheat): 11 Dec.

Potatoes: Spring-tine cultivated: 12 Mar, 1985. NPK Mg applied: 3 Apr. Rotary harrowed, potatoes planted: 9 Apr. Rotary ridged: 30 Apr. Weedkillers applied: 16 May. Mancozeb applied: 20 June, 3 July, 23 July, 6 Aug with pirimicarb on the second and third occasion. Fentin hydroxide applied: 21 Aug, 11 Sept. Haulm mechanically destroyed: 17 Sept. Lifted: 16 Oct.

S. barley: NPK applied: 11 Mar, 1985. Spring-tine cultivated, rotary harrowed, seed sown: 12 Mar. Weedkillers and fungicide applied: 16 May. Combine harvested: 21 Aug.

S. beans: Phorate applied: 11 Mar, 1985. Spring-tine cultivated, rotary harrowed, seed sown: 12 Mar. Weedkiller applied: 13 Mar. Benomyl and pirimicarb applied: 8 July. Combine harvested: 25 Sept.

85/R/RN/17

Cultivations, etc.:- (continued)

W.wheat: PK applied: 11 Dec, 1984. Seed sown, spring-tine
cultivated: 13 Dec. N applied: 16 Apr, 1985. Weedkillers and
tridemorph applied: 16 May. Propiconazole, carbendazim and maneb
applied: 3 July. Insecticide applied: 10 July. Combine
harvested: 6 Sept.

85/R/RN/17

SERIES I POTATOES

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	57.6
P6 K6 T	61.8
- - S	56.8
- - SA	63.0
- - SF	58.5
P2 - SA	60.0
P3 - SF	76.3
P4 - S	60.5
P5 - S	62.2
P5 - SF	60.2
P6 - S	65.1
- K2 SA	74.1
- K3 SF	66.5
- K4 S	57.1
- K5 S	60.2
- K5 SF	66.5
- K6 S	63.8
P1 K1 SA	63.0
P1 K3 SA	62.1
P2 K2 SA	62.8
P3 K1 SA	55.9
P3 K3 SA	68.2
P3 K4 SF	61.3
P4 K3 SF	66.4
P4 K4 S	63.0
P4 K5 S	61.3
P4 K5 SF	66.1
P4 K6 S	64.2
P5 K4 S	67.0
P5 K4 SF	66.0
P5 K5 S	64.3
P5 K6 S	58.5
P6 K4 S	61.7
P6 K5 S	61.2
P6 K6 S	64.0
MEAN	63.1

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
SED	3.07 MIN REP 2.51 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	5	2.17	3.5

85/R/RN/17

SERIES I POTATOES

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

TREATMNT	
- - -	96.2
P6 K6 T	95.8
- - S	96.6
- - SA	97.7
- - SF	95.4
P2 - SA	97.9
P3 - SF	96.8
P4 - S	97.1
P5 - S	96.3
P5 - SF	94.6
P6 - S	97.3
- K2 SA	97.4
- K3 SF	96.4
- K4 S	96.9
- K5 S	97.3
- K5 SF	94.9
- K6 S	97.3
P1 K1 SA	97.7
P1 K3 SA	96.5
P2 K2 SA	96.2
P3 K1 SA	96.4
P3 K3 SA	96.9
P3 K4 SF	95.5
P4 K3 SF	97.2
P4 K4 S	94.9
P4 K5 S	95.1
P4 K5 SF	96.8
P4 K6 S	97.8
P5 K4 S	96.6
P5 K4 SF	97.3
P5 K5 S	97.0
P5 K6 S	96.9
P6 K4 S	96.3
P6 K5 S	97.5
P6 K6 S	97.0
MEAN	96.6

PLOT AREA HARVESTED 0.00210

* SEDs APPLIES ONLY TO - - -, P6 K6 T, - - S, - - SR, P5 - S,
- K5 S, P4 K5 S AND P5 K4 S

	TREATMENT
MAX-MIN	- - S V ANY OF REMAINDER
MIN REP	ANY OF REMAINDER

85/R/RN/17

SERIES II SPRING BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	5.65
P6 K6 T	5.17
- - S	5.62
- - SA	5.43
- - SF	5.26
P2 - SA	5.57
P3 - SF	5.61
P4 - S	5.48
P5 - S	5.47
P5 - SF	4.82
P6 - S	5.55
- K2 SA	5.62
- K3 SF	5.40
- K4 S	5.58
- K5 S	5.70
- K5 SF	5.18
- K6 S	5.82
P1 K1 SA	5.45
P1 K3 SA	5.52
P2 K2 SA	5.34
P3 K1 SA	5.50
P3 K3 SA	5.33
P3 K4 SF	5.13
P4 K3 SF	5.03
P4 K4 S	5.31
P4 K5 S	5.35
P4 K5 SF	5.00
P4 K6 S	5.22
P5 K4 S	5.15
P5 K4 SF	5.10
P5 K5 S	5.09
P5 K6 S	5.51
P6 K4 S	4.92
P6 K5 S	5.82
P6 K6 S	4.98
MEAN	5.38

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
SED	0.277 MIN REP
	0.226 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	5	0.196	3.6
GRAIN MEAN DM%	80.2	PLOT AREA HARVESTED	0.00420

85/R/RN/17

SERIES III SPRING BEANS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	4.86
P6 K6 T	5.57
- - S	6.05
- - SA	6.20
- - SF	5.84
P2 - SA	6.37
P3 - SF	6.31
P4 - S	5.96
P5 - S	6.07
P5 - SF	5.69
P6 - S	5.64
- K2 SA	5.64
- K3 SF	5.32
- K4 S	4.44
- K5 S	5.78
- K5 SF	5.81
- K6 S	5.86
P1 K1 SA	6.40
P1 K3 SA	6.23
P2 K2 SA	6.08
P3 K1 SA	6.17
P3 K3 SA	5.56
P3 K4 SF	5.85
P4 K3 SF	6.08
P4 K4 S	5.65
P4 K5 S	5.58
P4 K5 SF	5.36
P4 K6 S	6.18
P5 K4 S	6.58
P5 K4 SF	5.58
P5 K5 S	5.71
P5 K6 S	5.62
P6 K4 S	4.87
P6 K5 S	5.64
P6 K6 S	5.60
MEAN	5.77

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.570 MIN REP
	0.465 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	5	0.403	7.0
GRAIN MEAN DM%	83.5	PLOT AREA HARVESTED	0.00386

85/R/RN/17

SERIES IV WINTER WHEAT

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	5.91
P6 K6 T	6.10
- - S	5.61
- - SA	5.44
- - SF	6.13
P2 - SA	5.15
P3 - SF	5.51
P4 - S	5.65
P5 - S	6.07
P5 - SF	4.89
P6 - S	5.45
- K2 SA	5.33
- K3 SF	6.04
- K4 S	5.80
- K5 S	5.94
- K5 SF	5.88
- K6 S	5.82
P1 K1 SA	5.67
P1 K3 SA	6.05
P2 K2 SA	5.25
P3 K1 SA	4.99
P3 K3 SA	6.46
P3 K4 SF	5.12
P4 K3 SF	6.21
P4 K4 S	5.87
P4 K5 S	5.79
P4 K5 SF	5.32
P4 K6 S	6.46
P5 K4 S	6.09
P5 K4 SF	5.82
P5 K5 S	6.04
P5 K6 S	5.78
P6 K4 S	4.92
P6 K5 S	6.12
P6 K6 S	5.80
MEAN	5.73

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.397 MIN REP
	0.324 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	5	0.281	4.9
GRAIN MEAN DM%	77.2	PLOT AREA HARVESTED	0.00420