

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1986

[Full Table of Content](#)



86/R/WW/6 N and Dcd - W. Wheat

Rothamsted Research

Rothamsted Research (1987) *86/R/WW/6 N and Dcd - W. Wheat* ; Yields Of The Field Experiments 1986, pp 194 - 196 - DOI: <https://doi.org/10.23637/ERADOC-1-36>

86/R/WW/6

WINTER WHEAT

N AND DCD

Object: To study the effects of a nitrification inhibitor in combination with different rates and timings of N on yield - New Zealand.

Sponsors: G.A. Rodgers, F.V. Widdowson.

Design: 2 randomised blocks of 26 plots.

Whole plot dimensions: 3.0 x 11.0.

Treatments: All combinations of:-

1. N INHIB Nitrification inhibitor added to nitrogen fertilizer:

NONE	None
DICYANDI	Dicyandiamide at 16 kg, divided equally between applications

2. N TIME Time and division of nitrogen fertilizer:

1111	Quarter of N on each of 13 Mar, 1986, 4 Apr, 28 Apr, 16 May
22--	Half of N on each of 13 Mar, 4 Apr
2-2-	Half of N on each of 13 Mar, 28 Apr
-22-	Half of N on each of 4 Apr, 28 Apr
4---	All of N on 13 Mar
--4-	All of N on 28 Apr

3. N RATE Amount of nitrogen fertilizer applied (kg N):

160	160
240	240

plus one extra treatment

EXTRA

NONE	No nitrogen fertilizer or inhibitor (duplicated)
------	--

NOTE: Nitrogen was applied as a mixture of urea and ammonium nitrate (28% N).

Basal applications: Weedkillers: Paraquat at 0.60 kg ion in 200 l. Clopyralid at 0.07 kg, bromoxynil octanoate at 0.34 kg, mecoprop at 2.5 kg and isoproturon at 2.1 kg in 200 l applied with the prochloraz and carbendazim. Fungicides: Prochloraz at 0.40 kg and carbendazim at 0.15 kg. Propiconazole at 0.12 kg in 200 l.

Seed: Avalon, sown at 190 kg.

86/R/WW/6

Cultivations, etc.:— Heavy spring-tine cultivated: 30 Aug, 1985.
Paraquat applied: 24 Sept. Heavy spring-tine cultivated, rotary
harrowed, seed sown: 1 Oct. Remaining weedkillers with prochloraz
and carbendazim applied: 30 Apr, 1986. Propiconazole applied:
26 June. Combine harvested: 22 Aug. Previous crops: W. wheat 1984,
w. oats 1985.

NOTE: A green crop sample was taken at the end of June to measure dry
matter, ear numbers and N content. The N content of grain was
determined.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

N TIME	1111	22--	2-2-	-22-	4---	--4-	MEAN
N INHIB							
NONE	9.26	9.14	8.90	9.35	8.73	9.34	9.12
DICYANDI	8.89	9.01	9.16	9.16	8.94	9.19	9.06
MEAN	9.08	9.08	9.03	9.26	8.83	9.27	9.09
N RATE	160	240	MEAN				
N INHIB							
NONE	9.00	9.24	9.12				
DICYANDI	8.89	9.24	9.06				
MEAN	8.94	9.24	9.09				
N RATE	160	240	MEAN				
N TIME							
1111	8.96	9.20	9.08				
22--	8.94	9.21	9.08				
2-2-	8.82	9.23	9.03				
-22-	9.03	9.48	9.26				
4---	8.71	8.96	8.83				
--4-	9.18	9.35	9.27				
MEAN	8.94	9.24	9.09				
N INHIB	N RATE	160	240				
NONE	N TIME						
	1111	9.11	9.41				
	22--	8.99	9.30				
	2-2-	8.96	8.83				
	-22-	9.19	9.51				
	4---	8.57	8.88				
	--4-	9.16	9.53				
DICYANDI	1111	8.80	8.98				
	22--	8.89	9.13				
	2-2-	8.69	9.64				
	-22-	8.87	9.45				
	4---	8.84	9.04				
	--4-	9.21	9.18				
NONE	6.04						
GRAND MEAN	8.86						

86/R/WW/6

GRAIN TONNES/HECTARE

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	N INHIB	N TIME	N RATE	N INHIB N TIME
SED	0.097	0.168	0.097	0.238
TABLE	N INHIB N RATE	N TIME N RATE	N INHIB N TIME N RATE	
SED	0.137	0.238	0.336	

SED FOR COMPARING NONE WITH ANY ITEM IN
N INHIB.N TIME.N RATE TABLE IS 0.291

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	26	0.336	3.8

GRAIN MEAN DM% 76.3

PLOT AREA HARVESTED 0.00251