

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1986

[Full Table of Content](#)



86/R/RA/9 Forms and Times of N - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1987) *86/R/RA/9 Forms and Times of N - W. Oilseed Rape ; Yields Of The Field Experiments 1986*, pp 285 - 286 - DOI: <https://doi.org/10.23637/ERADOC-1-36>

86/R/RA/9

WINTER OILSEED RAPE

FORMS AND TIMES OF N

Object: To compare the effects of single and divided dressings of urea and 'Nitro-Chalk' on the yield of w. oilseed rape - Little Hoos.

Sponsor: R.J. Darby.

Design: 3 randomised blocks of 15 plots.

Whole plot dimensions: 4.0 x 13.0.

Treatments: All combinations of:-

1. N FORM Forms of nitrogen fertilizer:

AMM NIT	Ammonium nitrate, (as 'Nitro-Chalk' 26% N)
UREA	Prilled urea

2. N TIME Times of applying a total dressing of 200 kg N:

4 - - -	All on 12 Mar, 1986
3 1 - -	Three quarters on 12 Mar, one quarter on 26 Mar
3 - 1 -	Three quarters on 12 Mar, one quarter on 24 Apr
2 2 - -	Half on 12 Mar, half on 26 Mar
2 - 2 -	Half on 12 Mar, half on 24 Apr
2 1 1 -	Half on 12 Mar, quarter on 26 Mar, quarter on 24 Apr
1 1 1 1	One quarter on 12 Mar and 26 Mar and 24 Apr and 7 May

plus one extra treatment

EXTRA

NONE No nitrogen fertilizer

Basal applications: Weedkillers: Fluazifop-P-butyl at 0.19 kg in 200 l with a wetting agent ('Agral' at 0.20 l). Clopyralid and propyzamide (as 'Matrikerb' at 1.6 kg) in 500 l. Fungicides: Iprodione at 0.50 kg in 500 l. Insecticide: Deltamethrin at 0.0062 kg in 500 l. Desiccant: Diquat at 0.60 kg in 500 l with a wetting agent ('Agral' at 0.50 l).

Seed: Mikado, dressed gamma HCH, thiram and fenpropimorph, sown at 8.0 kg.

Cultivations, etc.: Rotary grubbed: 17 Aug, 1985. Discd, spring-tine cultivated, seed sown: 22 Aug. Fluazifop-P-butyl with wetting agent applied: 27 Sept. Insecticide applied: 31 Oct. 'Matrikerb' applied: 19 Dec. Fungicide applied: 25 June, 1986. Desiccant with wetting agent applied: 24 July. Combine harvested: 3 Aug. Previous crops: W. wheat 1984, w. barley 1985.

NOTE: Incidence of light leaf spot and downy mildew on pods was assessed at the end of July.

86/R/RA/9

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** TABLES OF MEANS *****

	N FORM N TIME	AMM NIT	UREA	MEAN
4	- - -	4.08	3.75	3.92
3	1 - -	4.12	3.76	3.94
3	- 1 -	4.01	3.82	3.91
2	2 - -	4.09	3.96	4.03
2	- 2 -	3.77	3.80	3.79
2	1 1 -	3.94	3.83	3.88
1	1 1 1	3.96	3.92	3.94
	MEAN	3.99	3.84	3.91

EXTRA NONE 1.67

GRAND MEAN 3.77

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	N FORM	N TIME	N FORM N TIME & NONE
-----	-----	-----	-----
SED	0.059	0.110	0.156

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	28	0.191	5.1
GRAIN MEAN DM%	89.2		
PLOT AREA HARVESTED	0.00300		