

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ROTHAMSTED RESEARCH
Rothamsted Experimental Station
Harpenden
Herts
AL5 2JQ
UK
Tel: 0454 616161
Fax: 0454 616162
Email: [Rothamsted@ Rothamsted.ac.uk](mailto: Rothamsted@ Rothamsted.ac.uk)
Web: [Rothamsted.ac.uk](http:// Rothamsted.ac.uk)

87/R/B/6 and 87/W/B/6 Varieties and N - S. Barley

Rothamsted Research

Rothamsted Research (1988) *87/R/B/6 and 87/W/B/6 Varieties and N - S. Barley* ; Yields Of The Field Experiments 1987, pp 189 - 191 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/B/6 and 87/W/B/6

SPRING BARLEY

VARIETIES AND N

Object: To study the yields of some of the newer varieties of s. barley at two rates of nitrogen - Rothamsted (R), Highfield IV and Woburn (W), Lansome II.

Sponsor: R. Moffitt.

Design: 3 randomised blocks of 2 plots split into 10.

Sub-plot dimensions: (R) 3.0 x 10.0. (W) 4.0 x 9.0.

Treatments: All combinations of:-

Whole plots

1. N Nitrogen fertilizer:

(R)	(W)	
125	80	125 kg N on 20 Mar, 1987 (R), 80 kg N on 5 May (W)
125+46	120	125 kg N on 20 Mar plus 46 kg N on 29 Apr (R), 120 kg N on 5 May (W)

Sub plots

2. VARIETY Varieties:

BLENHEIM	Blenheim
CAMEO	Cameo
CORNICHE	Corniche
DIGGER	Digger
DOUBLET	Doublet
KLAXON	Klaxon
KLAXON B	Klaxon with 'Baytan' seed dressing
NATASHA	Natasha
REGATTA	Regatta
TRIUMPH	Triumph

NOTE: Nitrogen fertilizer was applied as 'Nitram'.

Basal applications:

Highfield IV (R): Weedkillers: Clopyralid at 0.07 kg, bromoxynil at 0.34 kg with mecoprop at 2.5 kg in 200 l. Fungicide: Tridemorph at 0.52 kg in 200 l.

Lansome II (W): Fungicides: Tridemorph at 0.19 kg with propiconazole at 0.12 kg in 200 l. Desiccant: Diquat at 0.60 kg ion applied with a wetting agent ('Agral' at 0.10 l) in 200 l.

Seed: Highfield IV (R), and Lansome II (W): Sown at 160 kg.

Cultivations, etc.:-

Highfield IV (R): Ploughed: 5 Nov, 1986. Spring-tine cultivated, rotary harrowed, seed sown, harrowed: 20 Mar, 1987. Rolled: 21 Mar. Weedkillers applied: 6 May. Fungicide applied: 23 June. Combine harvested: 20 Aug. Previous crops: Potatoes 1985, w. wheat 1986.

87/R/B/6 and 87/W/B/6

Cultivations, etc.:-

Lansome II (W): Deep-tine cultivated: 30 Jan, 1987. Spike harrowed with crumbler attached, seed sown: 30 Mar. Fungicides applied: 3 July. Desiccant applied: 21 Aug. Combine harvested: 10 Sept. Previous crops: W. oats 1985, potatoes 1986.

87/R/B/6 HIGHFIELD IV (R)

GRAIN TONNES/HECTARE

***** Tables of means *****

N	125	125+46	Mean
VARIETY			
BLENHEIM	6.20	6.59	6.39
CAMEO	6.99	7.63	7.31
CORNICHE	6.61	6.62	6.62
DIGGER	5.99	7.96	6.97
DOUBLET	6.81	7.52	7.17
KLAXON	6.24	6.55	6.39
KLAXON B	6.10	7.19	6.64
NATASHA	6.67	6.85	6.76
REGATTA	7.55	7.58	7.57
TRIUMPH	5.67	6.67	6.17
Mean	6.48	7.12	6.80

*** Standard errors of differences of means ***

Table	VARIETY	N*
s.e.d.	0.270	0.382

* Within the same level of N only

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP.SP	36	0.467	6.9

GRAIN MEAN DM% 86.7

SUB PLOT AREA HARVESTED 0.00204

87/W/B/6 LANSOME II (W)

GRAIN TONNES/HECTARE

***** Tables of means *****

N	80	120	Mean
VARIETY			
BLENHEIM	4.92	4.78	4.85
CAMEO	5.00	4.22	4.61
CORNICHE	5.17	4.63	4.90
DIGGER	5.94	6.68	6.31
DOUBLET	4.85	5.23	5.04
KLAXON	5.39	4.44	4.91
KLAXON B	5.02	4.86	4.94
NATASHA	4.57	4.22	4.39
REGATTA	5.43	5.17	5.30
TRIUMPH	5.26	4.03	4.64
Mean	5.15	4.83	4.99

*** Standard errors of differences of means ***

Table	VARIETY	N*
s.e.d.	0.236	0.334

* Within the same level of N only

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP.SP	36	0.409	8.2
GRAIN MEAN DM%	85.5		
SUB PLOT AREA HARVESTED	0.00247		