

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

87/R/RA/2
Seed Rates and Row Spacings - W. Oilseed Rape
Yields of the Field Experiments 1987
pp 213 - 214

87/R/RA/2 Seed Rates and Row Spacings - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1988) *87/R/RA/2 Seed Rates and Row Spacings - W. Oilseed Rape* ; Yields Of The Field Experiments 1987, pp 213 - 214 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/RA/2

WINTER OILSEED RAPE

SEED RATES AND ROW SPACINGS

Object: To compare c.v. Ariana on a range of row-widths and seed rates - Great Knott II.

Sponsor: D.P. Yeoman.

Design: 3 randomised blocks of 11 plots.

Whole plot dimensions: 3.0 x 15.0.

Treatments: All combinations of:-

1. SEEDRATE Seed rates:

4 KG
6 KG
8 KG

2. ROWSPACE Row spacings:

17.5 CM
35 CM
52.5 CM

plus two extra treatments, sown at 2 kg seed rate:-

EXTRA Row spacings:

2 KG 35 35 cm
2 KG 52.5 52.5 cm

Basal applications: Manures: 'Nitram' at 140 kg and later at 720 kg.
Weedkillers: Sodium trichloroacetate at 16 kg in 200 l. Clopyralid and propyzamide (as 'Matrikerb' at 1.6 kg) in 500 l. Fungicides: Prochloraz at 0.50 kg in 200 l. Iprodione at 0.50 kg in 200 l. Insecticides: Azinphos methyl at 0.40 kg and demeton-S-methyl sulphone at 0.12 kg in 300 l. Triazophos at 0.42 l in 200 l. Bird repellent: 'Hoppit' at 3.0 l in 220 l. Desiccant: Diquat at 0.60 kg ion with a wetting agent ('Enhance' at 0.50 l) in 500 l.

Seed: Ariana, dressed iprodione, gamma HCH and captan.

Cultivations, etc.: Heavy spring-tine cultivated, cultivated with rotary grubber: 12 Aug, 1986. First N applied: 15 Aug. Rotary harrowed: 31 Aug. Sodium trichloroacetate applied, harrowed: 1 Sept. Seed sown: 3 Sept. Remaining weedkillers applied: 20 Nov. Bird repellent applied: 12 Dec. Second N applied: 18 Feb, 1987. Prochloraz applied: 22 Apr. Azinphos methyl and demeton-S-methyl sulphone applied: 29 Apr. Iprodione and triazophos applied: 15 June. Desiccant with wetting agent applied: 28 July. Combine harvested: 5 Aug. Previous crops: S. wheat 1985, w. barley 1986.

NOTE: Plant counts were made at establishment and again in spring.

87/R/RA/2

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

ROWSPACE	17.5 CM	35 CM	52.5 CM	Mean
SEEDRATE				
4 KG	3.45	3.36	3.48	3.43
6 KG	3.17	3.13	2.90	3.07
8 KG	3.02	3.02	2.94	2.99
Mean	3.21	3.17	3.11	3.16

EXTRA	2 KG	35	2 KG	52.5	Mean
		3.61		3.39	3.50

GRAND MEAN 3.22

*** Standard errors of differences of means ***

Table	EXTRA	SEEDRATE	ROWSPACE	SEEDRATE ROWSPACE
s.e.d.	0.229	0.132	0.132	0.229

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	20	0.280	8.7

GRAIN MEAN DM% 71.3

PLOT AREA HARVESTED 0.00345