

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

87/R/RA/4 Growth Regulators and Fungicides - W. Oilseed Rape

Yields of the Field Experiments 1987, pp 220 - 223 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/RA/4 Growth Regulators and Fungicides - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1988) *87/R/RA/4 Growth Regulators and Fungicides - W. Oilseed Rape ; Yields Of The Field Experiments 1987*, pp 220 - 223 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/RA/4

WINTER OILSEED RAPE

GROWTH REGULATORS AND FUNGICIDES

Object: To study the effects of a range of materials on the control of fungi and on the growth and the yield of w. oilseed rape - Black Horse I.

Sponsor: C.J. Rawlinson.

Design: Single replicate of 3 x 4 x 2 x 2.

Whole plot dimensions: 3.0 x 20.0.

Treatments: All combinations of:-

Whole plots

- | | |
|--------------|---|
| 1. FUNGICIDE | Fungicides: |
| NONE | None |
| PROCHLOR | Prochloraz at 0.50 kg |
| PROPICON | Propiconazole at 0.12 kg |
| 2. GRTH REG | Growth regulators: |
| NONE | None |
| MEPIQUAT | Mepiquat chloride at 0.915 kg + 2-chlorethyl-phosphonic acid at 0.465 kg |
| TRIAPENT | Triapenthenol (as 'U.K.244a' at 0.70 kg) with a wetting agent ('Cittowett' at 0.10 l) |
| BAS11100 | 'BAS 11100W' at 3.0 l |
| 3. VARIETY | Varieties: |
| ARIANA | |
| PRIMOR | |
| 4. APP TIME | Times of application: |
| AUTUMN | Autumn, on 27 Nov, 1986 |
| SPRING | Spring, on 15 Apr, 1987 |

NOTE: Treatment sprays were applied in 220 l.

Basal applications: Manures: (0:18:36) at 690 kg. 'Nitram' at 140 kg, and later at 800 kg. Weedkillers: Sodium trichloroacetate at 16 kg in 200 l. Metazachlor at 1.2 kg with fluazifop-P-butyl at 0.19 kg and a wetting agent ('Agral' at 0.20 l) in 200 l. Insecticides: Deltamethrin at 0.0062 kg in 200 l. Azinphos methyl at 0.40 kg and demeton-S-methyl sulphone at 0.12 kg in 300 l. Bird repellent: 'Hoppit' at 3.0 l in 220 l. Desiccant: Diquat at 0.60 kg ion with a wetting agent ('Enhance' at 0.50 l) in 500 l.

87/R/RA/4

Seed: Varieties, sown at 8.0 kg.

Cultivations, etc.:— Spring-tine cultivated: 8 Aug, 1986. PK applied: 11 Aug. Sodium trichloroacetate applied, N applied: 13 Aug. Seed sown: 2 Sept. Remaining weedkillers applied: 4 Oct. Deltamethrin applied: 11 Oct. Bird repellent applied: 12 Dec. Second N applied: 17 Feb, 1987. Remaining insecticides applied: 28 Apr. Desiccant with wetting agent applied: 28 July. Combine harvested: 3 Aug. Previous crops: W. wheat 1985, w. barley 1986.

NOTE: Plant heights and diseases were assessed throughout the season. Flowering dates were noted and plant population counts made at harvest. Growth analysis and plant structure measurements were made just before harvest, stubble stem population counts were made immediately after harvest. Components of yield were measured.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

GRTH REG FUNGICIDE	NONE	MEPIQUAT	TRIAPENT	BAS11100	Mean
NONE	2.71	2.84	3.10	2.75	2.85
PROCHLOR	3.19	2.98	3.24	3.20	3.15
PROPICON	2.82	2.78	3.17	3.04	2.95
Mean	2.90	2.86	3.17	2.99	2.98
VARIETY FUNGICIDE	ARIANA	PRIMOR	Mean		
NONE	3.24	2.45	2.85		
PROCHLOR	3.54	2.76	3.15		
PROPICON	3.25	2.66	2.95		
Mean	3.34	2.62	2.98		
VARIETY GRTH REG	ARIANA	PRIMOR	Mean		
NONE	3.36	2.45	2.90		
MEPIQUAT	3.30	2.43	2.86		
TRIAPENT	3.41	2.93	3.17		
BAS11100	3.30	2.69	2.99		
Mean	3.34	2.62	2.98		
APP TIME FUNGICIDE	AUTUMN	SPRING	Mean		
NONE	2.83	2.86	2.85		
PROCHLOR	3.19	3.11	3.15		
PROPICON	2.96	2.95	2.95		
Mean	2.99	2.97	2.98		

87/R/RA/4

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

APP TIME	AUTUMN	SPRING	Mean
GRTH REG			
NONE	2.88	2.93	2.90
MEPIQUAT	2.95	2.78	2.86
TRIAPENT	3.12	3.22	3.17
BAS11100	3.02	2.97	2.99
Mean	2.99	2.97	2.98
APP TIME	AUTUMN	SPRING	Mean
VARIETY			
ARIANA	3.39	3.29	3.34
PRIMOR	2.60	2.65	2.62
Mean	2.99	2.97	2.98
FUNGICIDE	VARIETY	ARIANA	PRIMOR
GRTH REG			
NONE	NONE	3.30	2.12
	MEPIQUAT	3.24	2.44
	TRIAPENT	3.41	2.78
	BAS11100	3.01	2.48
PROCHLOR	NONE	3.59	2.79
	MEPIQUAT	3.47	2.49
	TRIAPENT	3.55	2.93
	BAS11100	3.54	2.85
PROPICON	NONE	3.18	2.45
	MEPIQUAT	3.19	2.37
	TRIAPENT	3.28	3.07
	BAS11100	3.34	2.74
FUNGICIDE	APP TIME	AUTUMN	SPRING
GRTH REG			
NONE	NONE	2.68	2.74
	MEPIQUAT	2.81	2.86
	TRIAPENT	3.04	3.15
	BAS11100	2.79	2.70
PROCHLOR	NONE	3.28	3.10
	MEPIQUAT	3.23	2.72
	TRIAPENT	3.13	3.35
	BAS11100	3.14	3.26
PROPICON	NONE	2.69	2.94
	MEPIQUAT	2.81	2.75
	TRIAPENT	3.19	3.16
	BAS11100	3.13	2.94
FUNGICIDE	VARIETY	ARIANA	PRIMOR
APP TIME			
NONE	AUTUMN	3.21	2.45
	SPRING	3.27	2.46
PROCHLOR	AUTUMN	3.56	2.82
	SPRING	3.51	2.70
PROPICON	AUTUMN	3.39	2.52
	SPRING	3.11	2.79

87/R/RA/4

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

	VARIETY	ARIANA		PRIMOR	
GRTH REG	APP TIME	AUTUMN	SPRING	AUTUMN	SPRING
NONE		3.24	3.47	2.52	2.38
MEPIQUAT		3.38	3.22	2.53	2.33
TRIAPENT		3.43	3.39	2.81	3.04
BAS11100		3.50	3.09	2.54	2.84

*** Standard errors of differences of means ***

Table	FUNGCIDE	GRTH REG	VARIETY	APP TIME
s.e.d.	0.103	0.119	0.084	0.084

Table	FUNGCIDE	FUNGCIDE	GRTH REG	FUNGCIDE
	GRTH REG	VARIETY	VARIETY	APP TIME
s.e.d.	0.206	0.145	0.168	0.145

Table	GRTH REG	VARIETY	FUNGCIDE	FUNGCIDE
	APP TIME	APP TIME	GRTH REG	GRTH REG
			VARIETY	APP TIME
s.e.d.	0.168	0.119	0.291	0.291

Table	FUNGCIDE	GRTH REG
	VARIETY	VARIETY
	APP TIME	APP TIME
s.e.d.	0.206	0.237

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
WP	6	0.291	9.7

GRAIN MEAN DM% 83.6

PLOT AREA HARVESTED 0.00460