

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

87/R/CS/140 Chemical Reference Plots - S. Barley
Rothamsted Experimental Station
Harpenden
Herts
AL5 2JQ
UK
This report has been prepared by Rothamsted Research and is the property of Rothamsted Research. It is not to be distributed outside Rothamsted Research without the written permission of Rothamsted Research. The report is for the use of the recipient only and is not to be used for any other purpose without the written permission of Rothamsted Research.

87/R/CS/140 Chemical Reference Plots - S. Barley

Rothamsted Research

Rothamsted Research (1988) *87/R/CS/140 Chemical Reference Plots - S. Barley* ; Yields Of The Field Experiments 1987, pp 90 - 94 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/CS/140

CHEMICAL REFERENCE PLOTS

Object: To study the persistence in soil of agricultural chemicals applied annually, singly and in combination and their effects on soil microflora and on yield of continuous s. barley - Long Hoos V 3.

Sponsors: R.H. Bromilow, R. Macdonald.

The 14th year, s. barley.

For previous years see 74-86/R/CS/140.

Design: Single replicate of 32 plots.

Whole plot dimensions: 4.06 x 4.57.

Treatments, applied cumulatively except as stated:

All combinations of:-

- | | |
|----------------|--|
| 1. WEEDKLLR | Weedkiller in autumn: |
| NONE | None |
| GLYPHOS | Glyphosate at 1.4 kg to barley stubble each autumn from 1979 to 1984 at 0.72 kg in 1985 and at 0.54 kg in 1986 |
| 2. FUNGCIDE[1] | Fungicide in autumn: |
| NONE | None |
| TRIADIM | Triadimefon at 0.25 kg in autumn 1981, 1982, 1984, 1985 and 1986, 0.28 kg in autumn 1983 |
| 3. FUNGCIDE[2] | Fungicide in spring: |
| NONE | None |
| BENOMYL | Benomyl at 4 kg to seedbed |
| 4. INSECTCDE | Insecticide: |
| NONE | None |
| CHLORFEN | Chlorfenvinphos at 2 kg to the seedbed |
| 5. NEMACIDE | Nematicide: |
| NONE | None |
| ALDICARB | Aldicarb at 6 kg to the seedbed |

NOTE: Glyphosate and triadimefon were applied in 220 l on 22 Sept, 1986 and 13 Oct respectively. Other treatments were applied on 19 Mar, 1987.

87/R/CS/140

Basal applications: Manures: (0:18:36) at 1060 kg, 'Nitro-Chalk' at 550 kg. Weedkillers: Bentazone at 0.80 kg, dichlorprop at 1.08 kg, and MCPA at 0.64 kg in 220 l applied with the fungicide. Fungicide: Tridemorph at 0.52 kg.

Seed: Klaxon, seed not dressed, sown at 160 kg.

Cultivations, etc.: PK applied: 16 Sept, 1986. Rotary cultivated: 13 Oct. Ploughed: 26 Nov. Spring-tine cultivated, seedbed treatments applied, rotary harrowed, seed sown and N applied: 19 Mar, 1987. Weedkillers and fungicide applied: 6 May. Combine harvested: 21 Aug.

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNGICIDE[1]	NONE	TRIADIM	Mean
WEEDKLLR			
NONE	5.26	5.40	5.33
GLYPHOS	5.13	5.24	5.18
Mean	5.19	5.32	5.26
FUNGICIDE[2]	NONE	BENOMYL	Mean
WEEDKLLR			
NONE	5.42	5.24	5.33
GLYPHOS	5.26	5.10	5.18
Mean	5.34	5.17	5.26
FUNGICIDE[2]	NONE	BENOMYL	Mean
FUNGICIDE[1]			
NONE	5.35	5.04	5.19
TRIADIM	5.34	5.30	5.32
Mean	5.34	5.17	5.26
INSTCDE	NONE	CHLORFEN	Mean
WEEDKLLR			
NONE	5.31	5.35	5.33
GLYPHOS	5.26	5.10	5.18
Mean	5.29	5.23	5.26
INSTCDE	NONE	CHLORFEN	Mean
FUNGICIDE[1]			
NONE	5.31	5.08	5.19
TRIADIM	5.26	5.37	5.32
Mean	5.29	5.23	5.26

87/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

INSC TCDE	NONE	CHLORFEN	Mean
FUNGCIDE[2]			
NONE	5.33	5.36	5.34
BENOMYL	5.24	5.09	5.17
Mean	5.29	5.23	5.26
NEMACIDE	NONE	ALDICARB	Mean
WEEDKLLR			
NONE	5.45	5.21	5.33
GLYPHOS	5.21	5.15	5.18
Mean	5.33	5.18	5.26
NEMACIDE	NONE	ALDICARB	Mean
FUNGCIDE[1]			
NONE	5.23	5.16	5.19
TRIADIM	5.42	5.21	5.32
Mean	5.33	5.18	5.26
NEMACIDE	NONE	ALDICARB	Mean
FUNGCIDE[2]			
NONE	5.46	5.23	5.34
BENOMYL	5.20	5.14	5.17
Mean	5.33	5.18	5.26
NEMACIDE	NONE	ALDICARB	Mean
INSC TCDE			
NONE	5.38	5.19	5.29
CHLORFEN	5.27	5.18	5.23
Mean	5.33	5.18	5.26
WEEDKLLR	FUNGCIDE[2]	NONE	BENOMYL
NONE	FUNGCIDE[1]		
	NONE	5.41	5.11
	TRIADIM	5.44	5.36
GLYPHOS	NONE	5.29	4.96
	TRIADIM	5.23	5.24
WEEDKLLR	INSC TCDE	NONE	CHLORFEN
NONE	FUNGCIDE[1]		
	NONE	5.22	5.30
	TRIADIM	5.40	5.40
GLYPHOS	NONE	5.39	4.86
	TRIADIM	5.13	5.34

87/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

	INSCTCDE	NONE	CHLORFEN
WEEDKLLR	FUNGICIDE[2]		
NONE	NONE	5.40	5.45
	BENOMYL	5.22	5.26
GLYPHOS	NONE	5.26	5.27
	BENOMYL	5.27	4.93
	INSCTCDE	NONE	CHLORFEN
FUNGICIDE[1]	FUNGICIDE[2]		
NONE	NONE	5.39	5.31
	BENOMYL	5.23	4.85
TRIADIM	NONE	5.26	5.41
	BENOMYL	5.26	5.34
	NEMACIDE	NONE	ALDICARB
WEEDKLLR	FUNGICIDE[1]		
NONE	NONE	5.34	5.19
	TRIADIM	5.56	5.24
GLYPHOS	NONE	5.13	5.13
	TRIADIM	5.29	5.18
	NEMACIDE	NONE	ALDICARB
WEEDKLLR	FUNGICIDE[2]		
NONE	NONE	5.55	5.29
	BENOMYL	5.34	5.14
GLYPHOS	NONE	5.36	5.17
	BENOMYL	5.06	5.14
	NEMACIDE	NONE	ALDICARB
FUNGICIDE[1]	FUNGICIDE[2]		
NONE	NONE	5.50	5.20
	BENOMYL	4.96	5.11
TRIADIM	NONE	5.41	5.26
	BENOMYL	5.43	5.17
	NEMACIDE	NONE	ALDICARB
WEEDKLLR	INSCTCDE		
NONE	NONE	5.42	5.20
	CHLORFEN	5.47	5.23
GLYPHOS	NONE	5.35	5.18
	CHLORFEN	5.07	5.13
	NEMACIDE	NONE	ALDICARB
FUNGICIDE[1]	INSCTCDE		
NONE	NONE	5.35	5.26
	CHLORFEN	5.11	5.05
TRIADIM	NONE	5.42	5.11
	CHLORFEN	5.43	5.32

87/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

	NEMACIDE	NONE	ALDICARB
FUNGICIDE[2]	INSECTCDE		
NONE	NONE	5.43	5.23
	CHLORFEN	5.49	5.23
BENOMYL	NONE	5.34	5.15
	CHLORFEN	5.06	5.13

*** Standard errors of differences of means ***

Margins of two factor tables	0.151
Two factor tables	0.214
Three factor tables	0.303

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
WP	6	0.428	8.1

GRAIN MEAN DM% 87.6

PLOT AREA HARVESTED 0.00075