

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1988

[Full Table of Content](#)



88/R/EX/4 Exhaustion Land - S. Barley

Rothamsted Research

Rothamsted Research (1989) *88/R/EX/4 Exhaustion Land - S. Barley* ; Yields Of The Field Experiments 1988, pp 19 - 21 - DOI: <https://doi.org/10.23637/ERADOC-1-43>

88/R/EX/4

EXHAUSTION LAND

Object: To study the residual effects of manures applied 1876-1901, and of additional phosphate applied since 1986, on the yield of continuous s. barley - Hoosfield.

The 133rd year, s. barley.

For previous years see 'Details' 1967, 1973 and 74-87/R/EX/4.

Treatments: All combinations of:-

Whole plots

1. **OLD RES** Residues of manures applied annually 1876-1901:

 O None
 D Farmyard manure at 35 tonnes
 N 96 kg N as ammonium salts
 P 34 kg P as superphosphate
 NPKNAMG N and P as above plus 137 kg K as sulphate of potash,
 16 kg Na as sulphate of soda, 11 kg Mg as sulphate
 of magnesia
2. **P** Phosphate applied annually from 1986 as
 superphosphate until 1987, triple superphosphate
 since:

 O None
 P1 44 kg
 P2 87 kg
 P3 131 kg

plus all combinations of:-

1. **OLD RES** Residues of manures applied annually 1876-1901:

 O None
 D Farmyard manure at 35 tonnes
 N* 96 kg N as nitrate of soda
 PK 34 kg P as superphosphate, 137 kg K as sulphate of
 potash
 N*PK N, P and K as above
2. **N88** Nitrogen fertilizer (kg N) as 'Nitro-Chalk' until
 1985, as 'Nitram' since 1986 (basal until 1975, on
 a cyclic system since 1976):

 0
 48
 96
 144

NOTE: All plots of the combination OLD RES, P were given N at 144 kg as 'Nitram' and K at 83 kg as muriate of potash.

88/R/EX/4

Basal applications: Weedkillers: Glyphosate at 1.4 kg in 200 l. Mecoprop at 2.4 kg with clopyralid at 0.05 kg and bromoxynil at 0.24 kg in 200 l. Fungicides: Propiconazole at 0.12 kg and tridemorph at 0.25 kg in 200 l on two occasions.

Seed: Triumph, seed dressed triadimenol and fuberidazole, sown at 160 kg.

Cultivations, etc.:- Glyphosate applied: 17 Nov, 1987. P and K applied: 11 Dec. Ploughed: 14 Dec. Heavy spring-tine cultivated twice: 22 Feb, 1988, 23 Feb. Rotary harrowed, seed sown: 7 Mar. N applied: 13 Apr. Remaining weedkillers applied: 11 May. Fungicides applied: 17 May, 17 June. Combine harvested: 15 Aug.

PHOSPHATE PLOTS

GRAIN TONNES/HECTARE

***** Tables of means *****

	P	O	P1	P2	P3	Mean
OLD RES						
O		3.04	6.01	6.68	6.77	5.63
D		5.48	6.99	7.16	7.17	6.70
N		2.34	6.33	6.93	7.15	5.69
P		5.05	6.93	7.12	7.16	6.57
NPKNAMG		4.69	6.79	7.22	7.21	6.48
Mean		4.12	6.61	7.02	7.09	6.21

GRAIN MEAN DM% 84.2

STRAW TONNES/HECTARE

***** Tables of means *****

	P	O	P1	P2	P3	Mean
OLD RES						
O		1.47	3.23	3.04	3.56	2.83
D		2.68	3.74	3.75	3.26	3.36
N		1.27	3.34	3.94	3.25	2.95
P		2.78	4.00	4.45	3.91	3.78
NPKNAMG		2.23	3.28	3.19	3.50	3.05
Mean		2.09	3.52	3.67	3.50	3.19

STRAW MEAN DM% 87.4

PLOT AREA HARVESTED 0.00728

88/R/EX/4

NITROGEN PLOTS

GRAIN TONNES/HECTARE

***** Tables of means *****

N88	0	48	96	144	Mean
OLD RES					
O	1.06	1.41	2.03	1.71	1.55
D	2.09	3.71	4.83	4.96	3.90
N*	1.13	1.35	1.80	2.33	1.65
PK	1.48	2.95	4.30	4.55	3.32
N*PK	1.80	2.36	3.64	4.43	3.06
Mean	1.51	2.36	3.32	3.59	2.70

GRAIN MEAN DM% 81.8

NITROGEN PLOTS

STRAW TONNES/HECTARE

***** Tables of means *****

N88	0	48	96	144	Mean
OLD RES					
O	0.29	0.58	0.72	0.71	0.58
D	0.63	1.61	2.19	2.04	1.62
N*	0.36	0.56	0.72	1.07	0.68
PK	0.64	1.48	2.04	2.41	1.64
N*PK	0.65	0.85	1.35	2.07	1.23
Mean	0.52	1.02	1.40	1.66	1.15

STRAW MEAN DM% 88.0

PLOT AREA HARVESTED 0.00728