

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1990

[Full Table of Content](#)



90/R/WW/2 Control of Volunteers - W. Wheat

Rothamsted Research

Rothamsted Research (1991) *90/R/WW/2 Control of Volunteers - W. Wheat ; Yields Of The Field Experiments 1990*, pp 106 - 110 - DOI: <https://doi.org/10.23637/ERADOC-1-42>

90/R/WW/2

WINTER WHEAT

CONTROL OF VOLUNTEERS

Object: To compare methods of volunteer control in winter wheat after w. and s. barley - Delafield, Fosters Corner.

Sponsors: R. Moffitt, D.G. Christian.

Design: 3 replicates of 8 x 3 criss-cross.

Column plot dimensions: 6.0 x 20.0.

Treatments: All combinations of:-

Column plots

- | | |
|-------------|-------------------------------|
| 1. PRIMCULT | Primary cultivations: |
| NONE | None until just before sowing |
| DYNDRIVE | 'Bomford Dynadrive' |
| PLOUGH | Plough |
| TINE | Tine |
| 2. CULTDATE | Date of cultivations: |
| EARLY | 25 July, 1989 (Delafield) |
| | 17 Aug (Fosters Corner) |
| LATER | 15 Aug (Delafield) |
| | 7 Sept (Fosters Corner) |

Row plots

- | | |
|-------------|--|
| 3. PRSOWCON | Pre-sowing volunteer control: |
| GLYPHOS | Glyphosate at 0.27 kg in 200 l on 18 Oct, 1989
(Delafield), 1 Nov and 13 Nov (Fosters Corner) |
| PARAQUAT | Paraquat at 0.60 kg ion in 200 l on 18 Oct
(Delafield), 1 Nov (Fosters Corner) |
| NONE | None |

- NOTES:** (1) The 'Bomford Dynadrive' has a frame similar to a rotary cultivator but it has two rotating shafts containing flat, slightly twisted, spade-shaped tines. The front shaft drives the rear, it is fitted with twice the number of blades and rotates at about one third the speed of the rear shaft.
- (2) A 1 m strip of Plaisant w. barley was broadcast on the surface, by drill at 120 kg on one end of each plot. It was broadcast before any treatments were applied on 25 July, 1989 (Delafield) and 17 Aug (Fosters Corner).
- (3) All plots were heavy spring-tine cultivated on 1 Nov, 1989 (Delafield), 14 Nov (Fosters Corner) then rotary harrowed and seed sown by drill 7 Nov (Delafield), 15 Nov (Fosters Corner). The seed was spring-tine cultivated in, (Delafield), and harrowed in, (Fosters Corner), after sowing.
- (4) PRIMCULT TINE was heavy spring-tine cultivated twice.

90/R/WW/2

Basal applications: Manures: 'Nitram' at 120 kg and later at 460 kg.
Weedkillers: Chlorotoluron at 3.5 kg in 200 l. Bromoxynil at 0.25 kg, ioxynil at 0.25 kg and mecoprop at 2.0 kg with the prochloraz in 200 l (Delafield). Bromoxynil at 0.28 kg, ioxynil at 0.28 kg and mecoprop at 2.2 kg with the prochloraz in 200 l (Fosters Corner). Fungicides: Prochloraz at 0.40 kg. Chlorothalonil at 0.50 kg with propiconazole at 0.12 kg in 200 l (Delafield). Fenpropimorph at 0.75 kg in 200 l.

Seed: Mercia, sown at 180 kg.

Cultivations, etc.:- Chlorotoluron applied: 21 Nov, 1989 (Fosters Corner), 23 Nov (Delafield). N applied: 9 Mar, 1990 and 12 Apr. Bromoxynil, ioxynil and mecoprop with prochloraz applied: 10 Apr (Delafield), 25 Apr (Fosters Corner). Chlorothalonil applied with the propiconazole (Fosters Corner): 31 May. Fenpropimorph applied: 29 June. Combine harvested: 10 Aug (Delafield), 13 Aug (Fosters Corner). Previous crops: W. oilseed rape 1988, w. barley 1989 (Delafield); sunflowers 1988, s. barley 1989 (Fosters Corner).

NOTES: (1) Ears of volunteer plants were counted at anthesis of the sown crop.
(2) Percentage contamination of harvested grain by volunteer grain was measured.

90/R/WW/2 DELAFIELD W.WHEAT AFTER W.BARLEY

GRAIN TONNES/HECTARE

***** Tables of means *****

CULTDATE	EARLY	LATER	Mean	
PRIMCULT				
NONE	5.73	5.83	5.78	
DYNDRIVE	5.48	5.80	5.64	
PLOUGH	6.30	6.43	6.36	
TINE	5.84	5.89	5.86	
Mean	5.84	5.99	5.91	
PRISOWCON	GLYPHOS	PARAQUAT	NONE	Mean
PRIMCULT				
NONE	6.11	5.79	5.44	5.78
DYNDRIVE	5.85	5.94	5.14	5.64
PLOUGH	6.76	6.38	5.95	6.36
TINE	6.08	6.24	5.27	5.86
Mean	6.20	6.09	5.45	5.91
PRISOWCON	GLYPHOS	PARAQUAT	NONE	Mean
CULTDATE				
EARLY	6.10	5.92	5.50	5.84
LATER	6.30	6.25	5.40	5.99
Mean	6.20	6.09	5.45	5.91

90/R/WW/2 DELAFIELD W.WHEAT AFTER W.BARLEY

GRAIN TONNES/HECTARE

***** Tables of means *****

PRIMCULT	PRSOWCON CULTDATE	GLYPHOS	PARAQUAT	NONE
NONE	EARLY	6.18	5.57	5.44
	LATER	6.05	6.00	5.44
DYNDRIVE	EARLY	5.75	5.70	4.99
	LATER	5.94	6.17	5.29
PLOUGH	EARLY	6.61	6.22	6.08
	LATER	6.92	6.54	5.81
TINE	EARLY	5.85	6.18	5.48
	LATER	6.30	6.31	5.06

*** Standard errors of differences of means ***

PRIMCULT	CULTDATE	PRSOWCON	PRIMCULT CULTDATE
0.167	0.118	0.296	0.236
PRIMCULT PRSOWCON	CULTDATE PRSOWCON	PRIMCULT CULTDATE PRSOWCON	
0.369	0.325	0.445	
Except when comparing means with the same level(s) of			
PRIMCULT	0.354		
CULTDATE		0.317	
PRSOWCON	0.248	0.175	0.350
PRIMCULT.CULTDATE			0.419

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP1	4	0.363	6.1
BLOCK.WP2	14	0.288	4.9
BLOCK.WP1.WP2	28	0.388	6.6

GRAIN MEAN DM% 89.5

PLOT AREA HARVESTED 0.00075

90/R/WW/2 FOSTERS CORNER W.WHEAT AFTER S.BARLEY

GRAIN TONNES/HECTARE

***** Tables of means *****

CULTDATE	EARLY	LATER	Mean	
PRIMCULT				
NONE	5.61	5.75	5.68	
DYNDRIIVE	5.43	5.83	5.63	
PLOUGH	5.69	5.78	5.73	
TINE	5.56	5.65	5.60	
Mean	5.57	5.75	5.66	
PRROWCON	GLYPHOS	PARAQUAT	NONE	Mean
PRIMCULT				
NONE	5.74	5.43	5.87	5.68
DYNDRIIVE	5.59	5.52	5.77	5.63
PLOUGH	5.73	5.59	5.88	5.73
TINE	5.55	5.48	5.78	5.60
Mean	5.65	5.51	5.83	5.66
PRROWCON	GLYPHOS	PARAQUAT	NONE	Mean
CULTDATE				
EARLY	5.55	5.40	5.77	5.57
LATER	5.75	5.61	5.89	5.75
Mean	5.65	5.51	5.83	5.66
PRIMCULT	PRROWCON	GLYPHOS	PARAQUAT	NONE
NONE	EARLY	5.72	5.36	5.76
	LATER	5.76	5.50	5.98
DYNDRIIVE	EARLY	5.43	5.32	5.55
	LATER	5.76	5.73	5.99
PLOUGH	EARLY	5.60	5.47	5.99
	LATER	5.86	5.71	5.78
TINE	EARLY	5.46	5.46	5.75
	LATER	5.64	5.50	5.80

*** Standard errors of differences of means ***

	PRIMCULT	CULTDATE	PRROWCON	PRIMCULT CULTDATE
	0.083	0.059	0.236	0.118
	PRIMCULT PRROWCON	CULTDATE PRROWCON	PRIMCULT CULTDATE PRROWCON	
	0.265	0.246	0.300	
Except when comparing means with the same level(s) of				
PRIMCULT	0.264			
CULTDATE		0.246		
PRROWCON	0.140	0.099	0.197	
PRIMCULT.CULTDATE			0.298	

90/R/WW/2 FOSTERS CORNER W.WHEAT AFTER S.BARLEY

GRAIN TONNES/HECTARE

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP1	4	0.289	5.1
BLOCK.WP2	14	0.144	2.5
BLOCK.WP1.WP2	28	0.238	4.2

GRAIN MEAN DM% 89.1

PLOT AREA HARVESTED 0.00080