

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1990

[Full Table of Content](#)



Sunflowers

Rothamsted Research

Rothamsted Research (1991) *Sunflowers* ; Yields Of The Field Experiments 1990, pp 143 - 152 -
DOI: <https://doi.org/10.23637/ERADOC-1-42>

90/R/SU/1

SUNFLOWERS

VARIETIES AND SOWING DATES

Object: To study the effects of five sowing dates on the rates of vegetative and floral development, days to maturity, disease and yield of two varieties of sunflowers - Great Harpenden I.

Sponsors: V.J. Church, C.J. Rawlinson.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 3.5 x 10.0.

Treatments: All combinations of:-

1. VARIETY	Varieties:
SUNB 246	Sunbred 246
S 47	S47
2. SOWDATE	Sowing date:
15 MAR	15 March, 1990
5 APR	5 April
18 APR	18 April
30 APR	30 April
10 MAY	10 May

NOTE: Plots were netted against birds from sowing until harvest.

Basal applications: Manures: (13:13:21) at 380 kg. Weedkillers: Trifluralin at 1.1 kg in 220 l. Linuron at 0.50 kg in 220 l. Desiccant: Diquat at 0.60 kg ion in 220 l. Irrigation: 22 mm applied on two occasions.

Seed: Sown at 20 seeds per square metre.

Cultivations, etc.:- Ploughed: 1 Sept, 1989. Rolled: 4 Sept. Deep-tine cultivated: 24 Nov. Spring-tine cultivated, NPK applied: 12 Mar, 1990. Trifluralin applied and rotary cultivated: 14 Mar to **SOWDATE** 15 MAR, 4 Apr to **SOWDATE** 5 APR, 18 Apr to **SOWDATE** 18 APR, 30 Apr to **SOWDATE** 30 APR and 9 May to **SOWDATE** 10 MAY. Seed sown as **SOWDATE**. Rolled: 16 Mar to **SOWDATE** 15 MAR, 5 Apr to **SOWDATE** 5 APR, 18 Apr to **SOWDATE** 18 APR, 30 Apr to **SOWDATE** 30 APR and 10 May to **SOWDATE** 10 MAY. Linuron applied: 16 Mar to **SOWDATE** 15 MAR, 5 Apr to **SOWDATE** 5 APR, 19 Apr to **SOWDATE** 18 APR, 1 May to **SOWDATE** 30 APR and 10 May to **SOWDATE** 10 MAY. Irrigation applied: 3 and 25 May. Desiccant applied: 13 Aug to **VARIETY** S 47 **SOWDATE** 15 MAR, 22 Aug to **VARIETY** S 47 **SOWDATE** 5 APR and 18 APR, 29 Aug to **VARIETY** S 47 **SOWDATE** 30 APR and **VARIETY** SUNB 246 **SOWDATE** 15 MAR, 31 Aug to **VARIETY** S 47 **SOWDATE** 10 May, 5 Sept to **VARIETY** SUNB 246 **SOWDATE** 5 APR and 18 APR, 14 Sept to **VARIETY** SUNB 246 **SOWDATE** 30 APR and 26 Sept to **VARIETY** SUNB 246 **SOWDATE** 10 MAY. Hand harvested: 17 Aug **VARIETY** S 47 **SOWDATE** 15 MAR,

90/R/SU/1

Cultivations, etc.:-

29 Aug **VARIETY S 47 SOWDATE 5 APR** and 18 APR, 4 Sept **VARIETY S 47 SOWDATE 30 APR** and **VARIETY SUNB 246 SOWDATE 15 MAR**, 10 Sept **VARIETY S 47 SOWDATE 10 MAY**, 13 Sept **VARIETY SUNB 246 SOWDATE 5 APR** and 18 APR, 24 Sept **VARIETY SUNB 246 SOWDATE 30 APR** and 2 Oct **VARIETY SUNB 246 SOWDATE 10 MAY**. Previous crops: W. wheat 1988 and 1989.

NOTE: Plant heights and head diameters were measured and plants counted at maturity. Botrytis was assessed on ten occasions in late summer. Severely distorted and male sterile heads were counted.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

SOWDATE VARIETY	15 MAR	5 APR	18 APR	30 APR	10 MAY	Mean
SUNB 246	2.91	3.11	3.30	3.48	3.53	3.27
S 47	0.83	1.72	2.01	2.39	2.26	1.84
Mean	1.87	2.41	2.66	2.93	2.89	2.55

*** Standard errors of differences of means ***

VARIETY	SOWDATE	VARIETY SOWDATE
0.039	0.062	0.088

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	18	0.107	4.2

GRAIN MEAN DM% 76.5

PLOT AREA HARVESTED 0.00150

90/R/SU/2

SUNFLOWERS

ROW SPACINGS AND SEED RATES

Object: To study the effects of three row spacings and three seed rates on growth, disease and yield of sunflowers - Great Harpenden I.

Sponsors: V.J. Church, C.J. Rawlinson.

Design: 4 randomised blocks of 9 plots.

Whole plot dimensions: Wide and narrow rows: 3.0 x 10.0
Medium rows: 3.04 x 10.0.

Treatments: All combinations of:-

- | | |
|-------------|-------------------------|
| 1. ROW SPAC | Spacing between rows: |
| NARROW | 25 cm |
| MEDIUM | 38 cm |
| WIDE | 50 cm |
| 2. POPULATN | Seeds sown per hectare: |
| 80 | 80,000 |
| 120 | 120,000 |
| 160 | 160,000 |

NOTE: Plants were netted against birds from sowing until harvest.

Basal applications: Manures: (13:13:21) at 380 kg. Weedkillers: Trifluralin at 1.1 kg in 200 l. Linuron at 0.50 kg in 200 l. Desiccant: Diquat at 0.60 kg ion applied with a wetting agent, 'Vassgrow' at 0.22 l, in 220 l. Irrigation: 22 mm on each of two occasions.

Seed: Vincent.

Cultivations, etc.:- Ploughed: 1 Sept, 1989. Rolled: 4 Sept. Deep-tine cultivated: 24 Nov. Spring-tine cultivated, NPK applied: 12 Mar, 1990. Trifluralin applied, spring-tine cultivated twice: 24 Apr. Seed sown: 26 Apr. Linuron applied: 1 May. Irrigation applied: 3 and 25 May. Desiccant applied: 5 Sept. Hand harvested: 12 Sept. Previous crops: W. wheat 1988 and 1989.

NOTE: Plant heights and head diameters were measured and plants counted at maturity. Botrytis was assessed in late summer.

90/R/SU/2

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

POPULATN	80	120	160	Mean
ROW SPAC				
NARROW	3.17	3.55	3.03	3.25
MEDIUM	3.25	3.28	2.85	3.13
WIDE	3.43	3.26	3.18	3.29
Mean	3.28	3.37	3.02	3.22

*** Standard errors of differences of means ***

ROW SPAC	POPULATN	ROW SPAC POPULATN
0.127	0.127	0.220

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	24	0.311	9.7

GRAIN MEAN DM% 76.3

PLOT AREA HARVESTED NARROW	0.00125
PLOT AREA HARVESTED MEDIUM	0.00114
PLOT AREA HARVESTED WIDE	0.00150

90/R/SU/3

SUNFLOWERS

METHODS OF APPLYING FUNGICIDES

Object: To compare the effects of mist-blown and hydraulic applications of fungicide on the control of Botrytis and on the yield of sunflowers - Great Harpenden I.

Sponsors: V.J. Church, C.J. Rawlinson.

Design: 6 randomised blocks of 4 plots.

Whole plot dimensions: 3.5 x 10.0.

Treatments:

SPRAYER	Sprayers:
NONE	None
HYD	Standard hydraulic sprayer
HYD DLN	Hydraulic sprayer with drop leg nozzles
MIST BLO	Mist blower

NOTE: The sprayers applied carbendazim at 0.15 kg and prochloraz at 0.40 kg with vinclozolin at 0.75 kg in 440 l in HYD and MIST BLO and in 600 l in HYD DLN on 17 and 25 July, 1990.

NOTE: Plots were netted against birds from sowing until harvest.

Basal applications: Manures: (13:13:21) at 380 kg. Weedkillers: Trifluralin at 1.1 kg in 200 l. Linuron at 0.50 kg in 200 l. Desiccant: Diquat at 0.60 kg ion applied with a wetting agent, 'Vassgrow' at 0.22 l, in 220 l. Irrigation: 22 mm on each of two occasions and 25 mm on a third.

Seed: S47, sown at 16 seeds per square metre.

Cultivations, etc.:- Ploughed: 1 Sept, 1989. Rolled: 4 Sept. Deep-tine cultivated: 24 Nov. Spring-tine cultivated, NPK applied: 12 Mar, 1990. Trifluralin applied, spring-tine cultivated twice: 24 Apr. Seed sown, rolled: 25 Apr. Linuron applied: 1 May. Irrigation applied: 3, 25 May and 27 July. Desiccant applied: 29 Aug. Hand harvested: 4 Sept. Previous crops: W. wheat 1988 and 1989.

NOTE: Botrytis was assessed on four occasions during August. Plants were counted at harvest.

90/R/SU/3

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

SPRAYER	NONE	HYD	HYD DLN	MIST BLO	Mean
	2.37	2.26	2.28	2.51	2.35

*** Standard errors of differences of means ***

SPRAYER
0.075

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	15	0.129	5.5

GRAIN MEAN DM% 79.0

PLOT AREA HARVESTED 0.00150

90/R/SU/4

SUNFLOWERS

DIMETHIPIN AND MATURITY

Object: To study the effects of dimethipin and diquat on dates of maturity, amounts of disease and yield of sunflowers - Great Harpenden I.

Sponsors: V.J. Church, C.J. Rawlinson.

Design: 3 randomised blocks each containing 2 replicates of 3 treatments.

Whole plot dimensions: 3.5 x 10.0.

Treatments:

CHEMICAL	Chemical sprays:
NONE	None
DIMETHIP	Dimethipin at 0.50 kg in 220 l applied on 22 Aug, 1990
DIQUAT	Diquat at 0.60 kg ion applied with a wetting agent, 'Vassgrow' at 0.22 l, in 220 l, applied on 5 Sept

NOTE: Plots were netted against birds from sowing until harvest.

Basal applications: Manures: (13:13:21) at 380 kg. Weedkillers: Trifluralin at 1.1 kg in 200 l. Linuron at 0.50 kg in 200 l. Irrigation: 22 mm on each of three occasions.

Seed: Vincent, sown at 16 seeds per square metre.

Cultivations, etc.:- Ploughed: 1 Sept, 1989. Rolled: 4 Sept. Deep-tine cultivated: 24 Nov. Spring-tine cultivated, NPK applied: 12 Mar, 1990. Trifluralin applied, spring-tine cultivated, rotary cultivated, seed sown, rolled: 24 Apr. Linuron applied: 1 May. Irrigation applied: 3, 10 and 25 May. Hand harvested: 10, 13 and 18 Sept for **CHEMICAL** DIMETHIP, DIQUAT and NONE respectively. Previous crops: W. wheat 1988 and 1989.

NOTE: Samples were assessed for seed moisture content during flowering. Botrytis was assessed three times during late summer. Plants were counted at harvest.

90/R/SU/4

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

CHEMICAL	NONE	DIMETHIP	DIQUAT	Mean
	2.71	2.51	2.67	2.63

*** Standard errors of differences of means ***

CHEMICAL
0.075

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	13	0.131	5.0

GRAIN MEAN DM% 80.8

PLOT AREA HARVESTED 0.00150

90/R/LN/1

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

FUNGCIDE	NONE	IP+PR+CM	Mean
INSCTCDE			
NONE	1.86	1.98	1.92
HC DE TR	1.85	2.07	1.96
Mean	1.86	2.03	1.94

*** Standard errors of differences of means ***

INSCTCDE	FUNGCIDE	INSCTCDE FUNGCIDE
0.037	0.037	0.052

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK	5	0.107	5.5
BLOCK.WP	15	0.091	4.7

GRAIN MEAN DM% 93.0

PLOT AREA HARVESTED 0.00276