

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1990

[Full Table of Content](#)



90/R/EX/4 Exhaustion Land - S. Barley

Rothamsted Research

Rothamsted Research (1991) *90/R/EX/4 Exhaustion Land - S. Barley* ; Yields Of The Field Experiments 1990, pp 19 - 21 - DOI: <https://doi.org/10.23637/ERADOC-1-42>

90/R/EX/4

EXHAUSTION LAND

Object: To study the residual effects of manures applied 1876-1901, and of additional phosphate applied since 1986, on the yield of continuous s. barley - Hoosfield.

The 135th year, s. barley.

For previous years see 'Details' 1967, 1973 and 74-89/R/EX/4.

Treatments: All combinations of:-

Whole plots

- | | |
|------------|--|
| 1. OLD RES | Residues of manures applied annually 1876-1901: |
| O | None |
| D | Farmyard manure at 35 tonnes |
| N | 96 kg N as ammonium salts |
| P | 34 kg P as superphosphate |
| NPKNAMG | N and P as above plus 137 kg K as sulphate of potash,
16 kg Na as sulphate of soda, 11 kg Mg as sulphate
of magnesia |
| 2. P | Phosphate applied annually from 1986 as
superphosphate until 1987, triple superphosphate
since: |
| O | None |
| P1 | 44 kg P |
| P2 | 87 kg P |
| P3 | 131 kg P |

plus all combinations of:-

- | | |
|------------|---|
| 1. OLD RES | Residues of manures applied annually 1876-1901: |
| O | None |
| D | Farmyard manure at 35 tonnes |
| N* | 96 kg N as nitrate of soda |
| PK | 34 kg P as superphosphate, 137 kg K as sulphate of
potash |
| N*PK | N, P and K as above |
| 2. N90 | Nitrogen fertilizer (kg N) as 'Nitro-Chalk' until
1985, as 'Nitram' since 1986 (basal until 1975, on
a cyclic system since 1976): |
| 0 | |
| 48 | |
| 96 | |
| 144 | |

NOTE: All plots of the combination OLD RES, P were given N at 144 kg as 'Nitram' and K at 83 kg as muriate of potash.

90/R/EX/4

Basal applications: Weedkillers: Glyphosate at 1.4 kg in 200 l.

Mecoprop at 1.6 kg, bromoxynil at 0.20 kg and ioxynil at 0.20 kg applied with the fungicide in 200 l. Fungicide: Fenpropimorph at 0.75 kg. Insecticide: Demeton-s-methyl at 0.24 kg in 200 l.

Seed: Triumph, seed dressed triadimenol and fuberidazole, sown at 160 kg.

Cultivations, etc.:- Glyphosate applied: 2 Oct, 1989. P and K applied:

1 Dec. Ploughed: 8 Dec. Spring-tine cultivated twice, rotary harrowed, seed sown: 9 Mar, 1990. N applied: 11 Apr. Insecticide applied: 10 May. Remaining weedkillers applied with the fungicide: 14 May. Combine harvested: 3 Aug.

PHOSPHATE PLOTS

GRAIN TONNES/HECTARE

***** Tables of means *****

P	O	P1	P2	P3	Mean
OLD RES					
O	1.91	3.79	3.90	4.42	3.50
D	3.42	4.35	4.66	4.10	4.13
N	1.97	3.96	4.53	4.45	3.73
P	2.96	4.19	4.17	4.07	3.85
NPKNAMG	2.90	3.96	4.62	4.29	3.94
Mean	2.63	4.05	4.38	4.26	3.83

GRAIN MEAN DM% 87.1

STRAW TONNES/HECTARE

***** Tables of means *****

P	O	P1	P2	P3	Mean
OLD RES					
O	0.36	1.30	1.31	1.30	1.07
D	1.20	1.40	1.60	1.59	1.45
N	0.46	1.31	1.41	1.40	1.15
P	0.93	1.68	1.40	1.77	1.45
NPKNAMG	1.11	1.48	1.79	1.49	1.47
Mean	0.81	1.44	1.50	1.51	1.32

STRAW MEAN DM% 93.2

PLOT AREA HARVESTED 0.00589

90/R/EX/4

NITROGEN PLOTS

GRAIN TONNES/HECTARE

***** Tables of means *****

N90	0	48	96	144	Mean
OLD RES					
O	1.18	1.39	1.23	0.82	1.16
D	1.92	2.40	2.26	2.33	2.23
N*	1.57	1.74	1.75	1.47	1.63
PK	2.26	2.13	2.68	2.48	2.39
N*PK	1.80	2.06	2.64	2.35	2.22
Mean	1.75	1.95	2.11	1.89	1.92

GRAIN MEAN DM% 86.1

STRAW TONNES/HECTARE

***** Tables of means *****

N90	0	48	96	144	Mean
OLD RES					
O	0.29	0.28	0.28	0.28	0.28
D	0.65	0.65	0.66	0.75	0.68
N*	0.57	0.56	0.47	0.37	0.49
PK	0.56	0.56	0.93	0.83	0.72
N*PK	0.47	0.75	0.75	0.74	0.68
Mean	0.51	0.56	0.62	0.59	0.57

STRAW MEAN DM% 93.7

PLOT AREA HARVESTED 0.00589