


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1990

[Full Table of Content](#)



90/R/CS/302 Eyespot Resistance to Mbc - W. Wheat

Rothamsted Research

Rothamsted Research (1991) *90/R/CS/302 Eyespot Resistance to Mbc - W. Wheat* ; Yields Of The Field Experiments 1990, pp 63 - 64 - DOI: <https://doi.org/10.23637/ERADOC-1-42>

90/R/CS/302

EYESPOT RESISTANCE TO MBC

Object: To study the development of resistance to MBC fungicides in eyespot and the ability of resistant strains to survive, spread and infect - Meadow.

Sponsor: G.L. Bateman.

The sixth year, w. wheat.

For previous years see 85-89/R/CS/302.

Design: 2 randomised blocks of 4 plots split into 6.

Whole plot dimensions: 12.0 x 24.0.

Treatments: All combinations of:-

Whole plots

1. **FUNGICIDE** Fungicides applied cumulatively to 1985, 1986, 1987, 1988 and 1989 treatments:

NONE	None
CARB	Carbendazim at 0.25 kg
PRO	Prochloraz at 0.40 kg
CARB+PRO	Carbendazim at 0.15 kg + prochloraz at 0.40 kg

Sub plots

2. **EYE INOC** Eyespot inoculum, applied in first year only:
- | | |
|----------|---|
| NATURAL | Natural background population (duplicated) |
| W 19R 1S | Inoculated with wheat strains in proportion 19 resistant to one sensitive |
| W 1R 19S | As above but one resistant to 19 sensitive |
| R 19R 1S | Inoculated with rye strains, 19 resistant to one sensitive |
| R 1R 19S | As above but one resistant to 19 sensitive |

NOTES: (1) Fungicide treatments were applied in 200 l on 24 Nov, 1989 and 30 Mar, 1990.
(2) The eyespot inoculum was colonised on oat seed and this was broadcast in October, 1984.

Basal applications: Manure: 'Nitram' at 580 kg. Weedkillers: Glyphosate at 0.27 kg in 200 l. Chlorotoluron at 3.0 kg with cyanazine at 0.75 kg in 200 l. Fluroxypyr at 0.15 kg with bromoxynil at 0.34 kg and clopyralid at 0.07 kg in 200 l. Insecticide: Deltamethrin at 6.2 g in 200 l.

Seed: Avalon, sown at 180 kg.

90/R/CS/302

Cultivations, etc.:- Heavy spring-tine cultivated: 22 Aug, 1989.

Glyphosate applied: 14 Sept. Ploughed: 22 Sept. Rotary harrowed twice, seed sown: 3 Oct. Chlorotoluron with cyanazine applied: 23 Nov. Deltamethrin applied: 22 Feb, 1990. N applied: 18 Apr. Fluroxypyr, bromoxynil and clopyralid applied: 30 Apr. Combine harvested: 10 Aug.

NOTE: Eyespot and sharp eyespot were assessed at fortnightly intervals from May - July on the EYE INOC NATURAL plots only.

GRAIN TONNES/HECTARE

***** Tables of means *****

EYE INOC FUNGICIDE	NATURAL	W 19R 1S	W 1R 19S	R 19R 1S	R 1R 19S	Mean
NONE	5.03	5.03	4.23	4.87	5.07	4.88
CARB	4.74	4.53	4.50	4.64	4.99	4.69
PRO	4.66	4.26	4.79	4.57	4.69	4.60
CARB+PRO	4.89	4.70	4.80	4.91	4.88	4.84
Mean	4.83	4.63	4.58	4.75	4.91	4.75

*** Standard errors of differences of means ***

EYE INOC	FUNGICIDE*
	EYE INOC
0.191	0.381 min.rep
0.165	0.330 max-min

EYE INOC
max-min NATURAL v any of the remainder
min.rep any of the remainder

* Within the same level of **FUNGICIDE** only

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP.SP	24	0.381	8.0

GRAIN MEAN DM% 90.1

SUB PLOT AREA HARVESTED 0.00138