


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1990

[Full Table of Content](#)



90/R/CS/311 Effects of Shallow Straw Incorporation - W. Wheat

Rothamsted Research

Rothamsted Research (1991) *90/R/CS/311 Effects of Shallow Straw Incorporation - W. Wheat* ; Yields Of The Field Experiments 1990, pp 69 - 73 - DOI: <https://doi.org/10.23637/ERADOC-1-42>

90/R/CS/311

EFFECTS OF SHALLOW STRAW INCORPORATION

Object: To study the effects of shallow straw incorporation on straw decomposition, toxin production, pests and diseases and on the establishment, growth and yield of winter wheat - West Barnfield I.

Sponsors: R.D. Prew, D.G. Christian, R.J. Gutteridge, E.T.G. Bacon, J.F. Jenkyn, B.R. Kerry, R. Moffitt, W. Powell, A.D. Todd.

The sixth year, w. wheat.

For previous years see 85-89/R/CS/311.

Design: Single replicate of 3 x a half replicate of 2 x 2 x 2 x 2 x 2.

Whole plot dimensions: 9.0 x 57.0.

Treatments: Combinations of:-

Whole plots

1. **STRAW** Treatments to straw of previous wheat:

BURNT	Burnt on 16 Aug, 1989
BALED	Baled and removed on 16 Aug
CHOPPED	Chopped on 16 Aug

2. **CULTTIME** Time of cultivation, to 10 cm depth:

EARLY	Cultivated by rotary grubber on 16 Aug, 1989
LATER	Cultivated by rotary grubber on 30 Aug

Sub plots

3. **AUTN RES** Residues of autumn N last applied autumn 1988, kg N per annum:

(0)
(50)

4. **FUNGICIDE** Fungicides:

O	None
FULL	Full programme:- Triadimefon at 0.12 kg and carbendazim at 0.25 kg in 200 l on 24 Nov, 1989. Prochloraz at 0.40 kg and carbendazim at 0.15 kg in 200 l on 9 Apr, 1990 Propiconazole at 0.125 kg in 200 l on 17 May Propiconazole at 0.125 kg with carbendazim at 0.25 kg and maneb at 1.6 kg in 200 l on 14 June

5. **INSECTCIDE** Insecticides:

O	None
CYP+PIR	Cypermethrin at 25 g in 200 l on 1 Nov, 1989 Pirimicarb at 0.14 kg in 200 l on 14 June

90/R/CS/311

6. MOLLICIDE

Molluscicide:

0 None
METHCARB Methiocarb at 0.22 kg on 6 Oct, 1989

NOTE: STRAW BURNT plots were disced the same day after burning.

Basal applications: Manures: 'Nitram' at 120 kg and later at 580 kg.
Weedkillers: Glyphosate at 0.27 kg in 200 l. Chlorotoluron at 3.0 kg with cyanazine at 0.75 kg in 200 l. Bromoxynil at 0.34 kg and clopyralid at 0.07 kg with fluroxypyr at 0.15 kg in 200 l.

Seed: Pastiche, sown at 200 kg.

Cultivations, etc.:- Glyphosate applied: 21 Sept, 1989. Rotary harrowed, seed sown: 5 Oct. Chlorotoluron with cyanazine applied: 10 Nov. First N applied: 2 Mar, 1990. Second N applied: 12 Apr. Remaining weedkillers applied: 3 May. Combine harvested: 11 Aug.

NOTE: Growth was measured and incidence of pests and diseases was assessed at intervals during the season. Ears of volunteers were counted prior to harvest and components of yield were measured.

GRAIN TONNES/HECTARE

***** Tables of means *****

CULTTIME	EARLY	LATER	Mean
STRAW			
BURNT	7.48	7.91	7.69
BALED	6.29	6.87	6.58
CHOPPED	5.55	7.01	6.28
Mean	6.44	7.27	6.85
AUTN RES	(0)	(50)	Mean
STRAW			
BURNT	7.73	7.65	7.69
BALED	6.50	6.66	6.58
CHOPPED	6.28	6.29	6.28
Mean	6.84	6.87	6.85
AUTN RES	(0)	(50)	Mean
CULTTIME			
EARLY	6.44	6.44	6.44
LATER	7.23	7.30	7.27
Mean	6.84	6.87	6.85
FUNGICIDE	0	FULL	Mean
STRAW			
BURNT	7.65	7.74	7.69
BALED	6.50	6.66	6.58
CHOPPED	6.27	6.30	6.28
Mean	6.81	6.90	6.85

90/R/CS/311

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNGCIDE	O	FULL	Mean
CULTTIME			
EARLY	6.40	6.48	6.44
LATER	7.21	7.32	7.27
Mean	6.81	6.90	6.85

FUNGCIDE	O	FULL	Mean
AUTN RES			
(0)	6.83	6.85	6.84
(50)	6.79	6.95	6.87
Mean	6.81	6.90	6.85

INSCDCDE	O	CYP+PIR	Mean
STRAW			
BURNT	7.47	7.92	7.69
BALED	6.50	6.67	6.58
CHOPPED	5.92	6.65	6.28
Mean	6.63	7.08	6.85

INSCDCDE	O	CYP+PIR	Mean
CULTTIME			
EARLY	6.11	6.77	6.44
LATER	7.14	7.39	7.27
Mean	6.63	7.08	6.85

INSCDCDE	O	CYP+PIR	Mean
AUTN RES			
(0)	6.63	7.05	6.84
(50)	6.63	7.11	6.87
Mean	6.63	7.08	6.85

INSCDCDE	O	CYP+PIR	Mean
FUNGCIDE			
O	6.63	6.98	6.81
FULL	6.62	7.18	6.90
Mean	6.63	7.08	6.85

MOLLCIDE	O	METHCARB	Mean
STRAW			
BURNT	7.68	7.71	7.69
BALED	6.55	6.62	6.58
CHOPPED	6.22	6.35	6.28
Mean	6.81	6.89	6.85

90/R/CS/311

GRAIN TONNES/HECTARE

***** Tables of means *****

MOLLCIDE CULTTIME	O	METHCARB	Mean
EARLY	6.37	6.51	6.44
LATER	7.26	7.27	7.27
Mean	6.81	6.89	6.85

MOLLCIDE AUTN RES	O	METHCARB	Mean
(0)	6.83	6.85	6.84
(50)	6.80	6.94	6.87
Mean	6.81	6.89	6.85

MOLLCIDE FUNGIDE	O	METHCARB	Mean
O	6.70	6.91	6.81
FULL	6.93	6.87	6.90
Mean	6.81	6.89	6.85

MOLLCIDE INSCDCDE	O	METHCARB	Mean
O	6.52	6.73	6.63
CYP+PIR	7.10	7.05	7.08
Mean	6.81	6.89	6.85

*** Standard errors of differences of means ***

AUTN RES	FUNGIDE	INSCDCDE	MOLLCIDE
0.072	0.072	0.072	0.072
STRAW*	CULTTIME*	STRAW*	CULTTIME*
AUTN RES	AUTN RES	FUNGIDE	FUNGIDE
0.125	0.102	0.125	0.102
AUTN RES	STRAW*	CULTTIME*	AUTN RES
FUNGIDE	INSCDCDE	INSCDCDE	INSCDCDE
0.102	0.125	0.102	0.102
FUNGIDE	STRAW*	CULTTIME*	AUTN RES
INSCDCDE	MOLLCIDE	MOLLCIDE	MOLLCIDE
0.102	0.125	0.102	0.102
FUNGIDE	INSCDCDE		
MOLLCIDE	MOLLCIDE		
0.102	0.102		

* Within the same level of STRAW, CULTTIME or STRAW.CULTTIME only

90/R/CS/311

GRAIN TONNES/HECTARE

**** Stratum standard errors and coefficients of variation ****

Stratum	d.f.	s.e.	cv%
WP.SP	20	0.250	3.6

GRAIN MEAN DM% 90.5

SUB PLOT AREA HARVESTED 0.00276