


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1991

[Full Table of Content](#)



Experiments - Classical

Rothamsted Research

Rothamsted Research (1992) *Experiments - Classical* ; Yields Of The Field Experiments 1991, pp 9 - 32 - DOI: <https://doi.org/10.23637/ERADOC-1-46>

91/R/BK/1

BROADBALK

Object: To study the effects of organic and inorganic manures on continuous w. wheat. From 1968 two three-year rotations were included: potatoes, beans, w. wheat and fallow, w. wheat, w. wheat. In 1979 the first rotation was changed to fallow, potatoes, w. wheat. In 1980 the second rotation reverted to continuous w. wheat. Since 1985 part of the second rotation has been added to the first to extend the rotation to fallow, potatoes, w. wheat, w. wheat, w. wheat.

The 148th year, w. wheat, fallow, potatoes.

For previous years see 'Details' 1967 and 1973, Station Report for 1966, pp. 229-231, Station Report for 1968, Part 2, and 74-90/R/BK/1.

Areas harvested:

Wheat:	Section	
	0	0.00311
	1	0.00572
	2, 4, 6 and 7	0.00473
	8 and 9	0.00497
Potatoes:	5	0.00348

Treatments:

Whole plots

PLOT

Fertilizers and organic manures:-

	Plot	Treatments until 1967	Treatments from 1968	Treatments from 1985
01DN4PK	01	-	D N2 P K	D N4 P K
21DN2	21	D	D N2	D N2
22D	22	D	D	D
030	03	None	None	None
05F	05	P K Na Mg	P K (Na) Mg	PK Mg
06N1F	06	N1 P K Na Mg	N1 P K (Na) Mg	N1 P K Mg
07N2F	07	N2 P K Na Mg	N2 P K (Na) Mg	N2 P K Mg
08N3F	08	N3 P K Na Mg	N3 P K (Na) Mg	N3 P K Mg
09N4F	09	N*1 P K Na Mg	N4 P K (Na) Mg	N4 P K Mg
10N2	10	N2	N2	N2
11N2P	11	N2 P	N2 P	N2 P
12N2PNa	12	N2 P Na	N2 P Na	N2 P Na
13N2PK	13	N2 P K	N2 P K	N2 P K
14N2PKMG	14	N2 P Mg	N2 P K Mg	N2 P K Mg
15N5F	15	N2 P K Na Mg	N3 P K (Na) Mg	N5 P K Mg
16N6F	16	N*2 P K Na Mg	N2 P K (Na) Mg	N6 P K Mg
17N0+3FH	17	N2 (A)	N2 1/2 (P K (Na) Mg)	N0+3 1/2 (PK Mg)+
18N1+3FH	18	P K Na Mg (A)	N2 1/2 (P K (Na) Mg)	N1+3 1/2 (PK Mg)+
19C	19	C	C	C
20NKMG	20	N2 K Na Mg	N2 K (Na) Mg	N2 K Mg

(A) Alternating

- N1,N2,N3,N4,N5,N6: 48, 96, 144, 192, 240, 288 kg N (as sulphate of ammonia until 1967, except N* which was nitrate of soda. All as 'Nitro-Chalk' in spring from 1968 to 1985, as 'Nitram' since 1986.)

F: P K (Na) Mg H: Half rate

70, 71, 72,
73, 74, 75,
and and and

W = w. wheat, P = potatoes, BE = s. beans, F = fallow

NOTES: (1) For a fuller record of treatments see 'Details' etc.
(2) From autumn 1975 to autumn 1986, chalk was applied at 2.9 t each autumn to all plots in sets of Sections on a three-year cycle. Year 1: Sections 1,2,3. Year 2: Sections 6,7,8,9. Year 3: Sections 0,4,5. Since autumn 1988 a five year cycle has been used. Year 1: Sections 1,3. Year 2: Sections 2,8. Year 3: Sections 7,9. Year 4: Sections 4,6. Year 5: Sections 0,5.

91/R/BK/1

Standard applications:

- W. wheat: Manure: Chalk at 2.9 t (to sections 7 and 9 only).
Weedkillers: Glyphosate at 1.4 kg in 200 l (except to sections 4 and 8). Diflufenican at 0.12 kg and isoproturon at 2.2 kg in 200 l (except to section 8). Glyphosate at 2.2 kg with a wetting agent, 'MAFF Adjuvant no. 0004' at 2.9 l, at 150 l (except to section 8).
Fungicides (except to section 6): Prochloraz at 0.40 kg with the growth regulator in 200 l. Propiconazole at 0.12 kg with chlorothalonil at 0.50 kg in 200 l. Fenpropimorph at 0.75 kg in 200 l. Growth regulator (except to section 6): Chlormequat chloride at 1.6 kg.
Potatoes: Weedkillers: Glyphosate at 1.4 kg in 200 l. Linuron at 1.6 kg in 200 l. Fungicides: Maneb at 0.96 kg and zinc oxide at 22 g in 200 l on three occasions, the first and third with a wetting agent, 'Bond' at 0.20 l, and the second occasion with pirimicarb. Mancozeb at 1.4 kg with a wetting agent, 'Bond' at 0.20 l, in 200 l. Fentin hydroxide at 0.27 kg with a wetting agent, 'Nu-film P' at 0.18 l, in 200 l. Insecticide: Pirimicarb at 0.14 kg.
Fallow: Weedkiller: Glyphosate at 2.2 kg with a wetting agent, 'MAFF Adjuvant no. 0004' at 2.9 l, in 150 l.

Seed: W. wheat: Apollo, dressed fonofos, sown at 180 kg.
Potatoes: Pentland Crown.

Cultivations, etc.:-

All Sections:

K, Na and Mg applied: 1 Oct, 1990. P applied: 2 Oct. FYM applied, ploughed and furrow pressed: 9 Oct. Rotary harrowed: 15 Oct.

Cropped Sections:

- W. wheat: Straw chopped (section 0): 14 Aug, 1990. Glyphosate alone applied (except to sections 4 and 8): 23 Aug. Chalk applied (sections 7 and 9): 27 Sept. Autumn N treatments applied: 2 Oct. Rotary harrowed, seed sown: 16 Oct. Diflufenican with isoproturon applied (except to section 8): 21 Nov. Spring N treatments applied: 9 Apr, 1991. Prochloraz with the growth regulator applied (except to section 6): 24 Apr. Propiconazole with chlorothalonil applied (except to section 6): 20 June. Fenpropimorph applied (except to section 6): 2 July. Glyphosate with the wetting agent applied (except to section 8): 12 Aug. Combine harvested: 25 Aug.
Potatoes: Glyphosate applied: 11 Sept, 1990. Deep-tine cultivated: 14 Dec. N treatments applied: 9 Apr, 1991. Heavy spring-tine cultivated: 16 Apr. Rotary harrowed, potatoes planted: 17 Apr. Rotary ridged: 9 May. Linuron applied: 21 May. Maneb and zinc oxide with the wetting agent applied: 1 July and 22 July. Maneb and zinc oxide with the insecticide applied: 10 July. Mancozeb with the wetting agent applied: 1 Aug. Fentin hydroxide with the wetting agent applied: 12 Aug. Haulm mechanically destroyed: 28 Aug. Lifted: 24 Sept.
Fallow: Deep-tine cultivated: 14 Dec, 1990. Heavy spring-tine cultivated: 25 Apr, 1991. Cultivated by rotary grubber: 21 June. Heavy spring-tine cultivated: 8 July. Glyphosate with wetting agent applied: 12 July.

91/R/BK/1 W.WHEAT

GRAIN TONNES/HECTARE

***** Tables of means *****

SECTION PLOT	4/W1	7/W2	2/W3	8/W3	6/W14	1/W25	9/W33	0/W40
01DN4PK	8.84	8.60	9.14	*	5.80	*	*	*
21DN2	9.38	8.65	8.84	3.31	7.18	8.98	9.10	8.74
22D	7.80	5.69	6.17	3.60	5.38	5.98	6.87	5.50
030	1.62	0.67	0.79	1.75	1.12	1.25	0.96	1.00
05F	1.50	0.52	0.85	2.10	1.17	1.29	1.09	1.51
06N1F	4.84	3.14	2.92	2.54	3.32	3.39	3.78	3.37
07N2F	6.92	5.42	5.07	3.27	5.75	6.29	6.19	5.82
08N3F	7.96	7.96	6.88	5.00	7.09	7.80	7.76	7.21
09N4F	8.35	8.43	7.39	5.63	6.95	7.65	8.21	7.34
10N2	5.63	4.85	3.29	1.76	3.54	3.14	4.00	3.11
11N2P	4.60	5.40	4.31	2.11	3.25	4.36	1.85	3.67
12N2PNA	5.32	5.67	3.92	2.32	4.31	2.85	2.22	4.51
13N2PK	6.74	4.95	4.56	3.28	5.15	5.42	6.09	5.34
14N2PKMG	6.72	5.27	4.69	2.86	5.14	5.77	6.20	5.67
15N5F	8.51	8.96	7.85	3.89	6.97	8.66	8.84	7.89
16N6F	8.26	8.50	8.21	3.77	6.28	8.16	8.22	7.32
17N0+3FH	8.27	7.40	6.39	3.28	6.36	7.22	7.30	6.91
18N1+3FH	8.38	7.81	7.25	4.33	6.62	7.28	7.38	7.42
19C	3.19	0.84	1.04	2.51	1.64	2.02	1.76	1.90
20NKMG	*	*	*	*	*	3.23	*	3.03

GRAIN MEAN DM% 82.7

STRAW TONNES/HECTARE

***** Tables of means *****

SECTION PLOT	4/W1	1/W25
01DN4PK	9.02	*
21DN2	9.11	8.67
22D	7.24	4.58
030	1.11	1.06
05F	0.90	0.92
06N1F	3.27	3.01
07N2F	5.54	4.75
08N3F	6.56	5.56
09N4F	6.60	5.90
10N2	3.04	3.78
11N2P	2.96	2.93
12N2PNA	3.37	1.68
13N2PK	5.57	4.38
14N2PKMG	5.86	3.98
15N5F	7.54	6.78
16N6F	7.05	6.34
17N0+3FH	6.68	5.95
18N1+3FH	6.83	6.46
19C	1.48	1.52
20NKMG	*	3.44

STRAW MEAN DM% 89.8

91/R/BK/1

POTATOES

***** Tables of means *****

PLOT	TOTAL TUBERS	% WARE
	TONNES/ HECTARE	3.81 CM (1.5 INCH) RIDDLE
01DN4PK	41.9	93.4
21DN2	45.7	95.6
22D	34.2	93.0
030	5.5	62.6
05F	10.6	78.0
06N1F	23.3	88.8
07N2F	29.2	88.8
08N3F	32.8	91.3
09N4F	29.7	91.4
10N2	8.7	78.2
11N2P	10.0	77.8
12N2PNA	9.8	73.2
13N2PK	18.3	75.9
14N2PKMG	31.1	90.6
15N5F	37.9	90.7
16N6F	35.3	93.2
17N3FH	22.1	89.5
18N3FH	27.1	92.5
19C	16.0	88.9

91/R/HB/2

HOOS BARLEY

Object: To study the effects of organic and inorganic manures on continuous s. barley. From 1968 to 1978 a rotation of potatoes, beans and s. barley was practised. The rotation was discontinued in 1979 and the experiment reverted to continuous s. barley.

The 140th year, s. barley.

For previous years see 'Details' 1967 and 1973, Station Report for 1966 and 74-90/R/HB/2.

Treatments: All combinations of:-

1. MANURE Fertilizers and organic manures:

	Form of N 1852-1966	Additional treatments 1852-1979	Changes since 1980
---	None	-	-
-P-	None	P	-
--K	None	K (Na) Mg	-
-PK	None	PK (Na) Mg	-
A--	A	-	-
AP-	A	P	-
A-K	A	K (Na) Mg	-
APK	A	PK (Na) Mg	-
N----	N	-	-
NP---	N	P	-
N-K--	N	K (Na) Mg	-
NPK--	N	PK (Na) Mg	-
N--S-	N	Si	Si omitted
NP-S-	N	P Si	"
N-KS-	N	K (Na) MgSi	"
NPKS-	N	PK (Na) MgSi	"
N---S	N	-	Si added
NP--S	N	P	"
N-K-S	N	K (Na) Mg	"
NPK-S	N	PK (Na) Mg	"
N--SS	N	Si	-
NP-SS	N	P Si	-
N-KSS	N	K (Na) MgSi	-
NPKSS	N	PK (Na) MgSi	-
C(--)	C	-	PKMg omitted
C(P-)	C	P	"
C(-K)	C	K (Na) Mg	"
C(PK)	C	PK (Na) Mg	"
D	None	D	-
(D)	(D)	-	-
(A)	(Ashes)	-	-
-	None	-	-

91/R/HB/2

Form of N: A, sulphate of ammonia: N, nitrate of soda - each to supply 48 kg N: C, castor meal to supply 96 kg N
 P: 35 kg P as triple superphosphate (triple superphosphate in 1974, 1988 and 1989, single superphosphate in other years)
 K: 90 kg K as sulphate of potash
 (Na): 16 kg Na as sulphate of soda until 1973
 Mg: 35 kg Mg, as kieserite every third year since 1974 (sulphate of magnesia annually until 1973)
 Si: Silicate of soda at 450 kg
 D: Farmyard manure at 35 tonnes. (D): until 1871 only
 (Ashes): Weed ash 1852-1916, furnace ash 1917-1932, none since

2. **N** Nitrogen fertilizer (kg N), as 'Nitro-Chalk', since 1968 (cumulative N applications until 1973, on a cyclic system since 1974):

0
 48
 96
 144

Plus extra plots testing all combinations of:-

1. **MANURE** Fertilizers other than magnesium:

551AN2PK	Plot 551 AN2PK
561--PK	Plot 561 --PK
571NN2--	Plot 571 NN2
581NN2--	Plot 581 NN2

N2: 96 kg N as 'Nitro-Chalk' since 1968. Other symbols as above.

2. **MAGNESIUM** Magnesium fertilizer (kg Mg) as kieserite every third year since 1974:

0
 35

NOTE: For a fuller record see 'Details' etc.

Basal applications: Weedkillers: Glyphosate at 1.4 kg in 200 l. Mecoprop at 1.1 kg, ioxynil at 0.20 kg, bromoxynil at 0.20 kg and linuron at 0.04 kg in 200 l. Fungicide: Tridemorph at 0.52 kg in 200 l. Insecticide: Pirimicarb at 0.14 kg in 200 l.

Seed: Triumph, dressed triadimenol and fuberidazole, sown at 130 kg.

Cultivations, etc.:- Glyphosate applied: 7 Nov, 1990. P applied: 21 Nov. K and silicate of soda applied: 28 Nov. FYM applied, ploughed: 3 Dec. Spring-tine cultivated, rotary harrowed, seed sown: 14 Mar, 1991. N applied: 12 Apr. Remaining weedkillers applied: 24 May. Fungicide applied: 4 June. Insecticide applied: 11 July. Combine harvested: 19 Aug.

91/R/HB/2

MAIN PLOTS

GRAIN TONNES/HECTARE

***** Tables of means *****

N	0	48	96	144	Mean
MANURE					
---	0.98	2.26	2.98	1.67	1.98
-P-	2.85	4.10	3.48	4.78	3.80
--K	2.52	4.02	4.62	4.14	3.82
-PK	2.66	4.58	5.74	6.21	4.80
A--	1.79	2.20	2.62	2.54	2.29
AP-	2.68	3.56	3.06	2.98	3.07
A-K	2.26	3.42	3.47	3.57	3.18
APK	2.67	4.43	5.39	5.82	4.57
N----	2.69	2.88	3.52	3.39	3.12
NP---	3.20	4.37	4.83	4.15	4.14
N-K--	2.69	3.82	3.83	3.56	3.48
NPK--	2.67	4.54	5.53	6.55	4.82
N--S-	2.67	4.63	3.50	3.49	3.57
NP-S-	3.33	4.45	4.59	5.48	4.46
N-KS-	2.85	4.17	4.94	5.13	4.27
NPKS-	2.99	4.80	5.64	6.50	4.98
N---S	2.83	3.23	3.76	4.32	3.54
NP--S	3.15	4.71	5.49	4.24	4.40
N-K-S	2.80	3.77	4.70	4.09	3.84
NPK-S	2.93	4.77	5.93	6.07	4.92
N--SS	2.62	3.22	4.09	4.22	3.54
NP-SS	3.21	4.45	4.42	5.02	4.27
N-KSS	2.99	4.23	4.93	4.56	4.18
NPKSS	1.93	5.62	6.07	6.20	4.96
C(--)	2.94	4.54	4.42	5.10	4.25
C(P-)	3.19	4.69	5.15	5.69	4.68
C(-K)	2.83	4.84	5.87	5.95	4.87
C(PK)	2.81	4.53	5.32	5.73	4.60
D	6.35	6.22	6.16	5.93	6.16
(D)	3.61	5.71	5.46	5.18	4.99
(A)	3.48	4.11	4.03	4.17	3.95
-	2.49	4.03	3.73	4.08	3.58
Mean	2.86	4.22	4.60	4.70	4.10

GRAIN MEAN DM% 87.8

91/R/HB/2

STRAW TONNES/HECTARE

***** Tables of means *****

N	0	48	96	144	Mean
MANURE					
---	0.42	0.91	1.34	0.70	0.84
-P-	1.14	1.91	1.73	2.53	1.83
--K	1.06	1.51	2.66	2.12	1.84
-PK	0.89	2.03	2.79	3.14	2.21
A--	0.55	0.75	1.30	1.20	0.95
AP-	0.93	1.70	1.91	1.74	1.57
A-K	0.87	1.58	1.75	1.59	1.45
APK	0.90	2.11	2.58	2.85	2.11
D	3.90	4.26	4.71	4.43	4.32
(D)	1.50	2.51	2.75	2.53	2.32
(A)	1.38	1.70	2.01	2.03	1.78
-	1.22	1.70	1.72	2.00	1.66
Mean	1.23	1.89	2.27	2.24	1.91

STRAW MEAN DM% 85.9

PLOT AREA HARVESTED 0.00154

EXTRA PLOTS

GRAIN TONNES/HECTARE

***** Tables of means *****

MANURE	551AN2PK	561--PK	571NN2--	581NN2--	Mean
MAGNESIUM					
0	5.11	1.64	5.24	3.24	3.81
35	5.50	1.53	5.11	3.26	3.85
Mean	5.31	1.59	5.18	3.25	3.83

GRAIN MEAN DM% 86.9

PLOT AREA HARVESTED 0.00329

91/R/WF/3

WHEAT AND FALLOW

Object: To study the effects of fallowing on unmanured w. wheat - Hoosfield.

The 136th year, w. wheat.

For previous years see 'Details' 1967, 1973 and 74-90/R/WF/3.

Whole plot dimensions: 9.0 x 211.

Treatments:

Each year there are two plots, one is sown to w. wheat, one is fallow; they alternate in successive years.

Seed: Apollo, dressed fonofos, sown at 180 kg.

Cultivations, etc.:-

Wheat plot: Rotary harrowed twice, seed sown: 17 Oct, 1990. Combine harvested: 25 Aug, 1991.

Fallow plot: Deep tine cultivated with vibrating tines 60 cm apart, 45 cm deep: 24 Sept, 1990. Ploughed, furrow pressed: 4 Oct. Heavy spring-tine cultivated: 25 Apr, 1991. Spring-tine cultivated: 14 June. Heavy spring-tine cultivated: 8 July.

GRAIN AND STRAW TONNES/HECTARE

	GRAIN	STRAW
YIELD	1.50	1.49
MEAN DM%	82.5	91.8
PLOT AREA HARVESTED	0.04309	

91/R/EX/4

EXHAUSTION LAND

Object: To study the residual effects of manures applied 1876-1901, and of additional phosphate applied since 1986, on the yield of continuous s. barley - Hoosfield.

The 136th year, s. barley.

For previous years see 'Details' 1967, 1973 and 74-90/R/EX/4.

Treatments: All combinations of:-

Whole plots

1. **OLD RES** Residues of manures applied annually 1876-1901:

O	None
D	Farmyard manure at 35 tonnes
N	96 kg N as ammonium salts
P	34 kg P as superphosphate
NPKNAMG	N and P as above plus 137 kg K as sulphate of potash, 16 kg Na as sulphate of soda, 11 kg Mg as sulphate of magnesia

2. **P** Phosphate applied annually from 1986 as
superphosphate until 1987, triple superphosphate
since:

O	None
P1	44 kg P
P2	87 kg P
P3	131 kg P

plus all combinations of:-

1. **OLD RES** Residues of manures applied annually 1876-1901:

O	None
D	Farmyard manure at 35 tonnes
N*	96 kg N as nitrate of soda
PK	34 kg P as superphosphate, 137 kg K as sulphate of potash
N*PK	N, P and K as above

2. **N91** Nitrogen fertilizer (kg N) as 'Nitro-Chalk' until
1985, as 'Nitram' since 1986 (basal until 1975, on
a cyclic system since 1976):

0
48
96
144

NOTE: All plots of the combination **OLD RES**, P were given N at 144 kg as 'Nitram' and K at 83 kg as muriate of potash.

91/R/EX/4

Basal applications: Weedkiller: Fluroxypyr at 0.15 kg with the fungicide in 200 l. Fungicide: Fenpropimorph at 0.38 kg.

Seed: Triumph, seed dressed triadimenol and fuberidazole, sown at 130 kg.

Cultivations, etc.:- P and K applied: 20 Nov, 1990. Ploughed: 23 Nov. Spring-tine cultivated: 25 Mar, 1991. Rotary harrowed, seed sown: 26 Mar. N applied: 18 Apr. Weedkiller with fungicide applied: 16 June. Combine harvested: 29 Aug.

PHOSPHATE PLOTS

GRAIN TONNES/HECTARE

***** Tables of means *****

	P	O	P1	P2	P3	Mean
OLD RES						
O		2.54	4.96	5.42	5.72	4.66
D		4.31	5.22	5.50	5.49	5.13
N		2.35	5.10	5.50	5.86	4.70
P		3.76	5.18	5.24	5.50	4.92
NPKNAMG		3.69	5.14	5.42	5.53	4.94
Mean		3.33	5.12	5.42	5.62	4.87

GRAIN MEAN DM% 85.1

STRAW TONNES/HECTARE

***** Tables of means *****

	P	O	P1	P2	P3	Mean
OLD RES						
O		1.82	3.69	4.14	3.81	3.36
D		3.05	3.72	3.71	3.53	3.50
N		1.73	3.59	3.70	3.79	3.20
P		2.40	3.60	3.61	3.62	3.31
NPKNAMG		2.50	3.45	3.74	3.70	3.35
Mean		2.30	3.61	3.78	3.69	3.34

STRAW MEAN DM% 92.4

PLOT AREA HARVESTED 0.00588

91/R/EX/4

NITROGEN PLOTS

GRAIN TONNES/HECTARE

***** Tables of means *****

	N91	0	48	96	144	Mean
OLD RES						
O		0.73	1.75	0.74	0.91	1.03
D		2.09	3.41	3.46	3.37	3.08
N*		0.96	2.13	1.15	1.14	1.34
PK		1.74	3.47	3.18	3.64	3.01
N*PK		1.71	3.41	2.99	2.88	2.75
Mean		1.44	2.83	2.30	2.39	2.24

GRAIN MEAN DM% 82.5

STRAW TONNES/HECTARE

***** Tables of means *****

	N91	0	48	96	144	Mean
OLD RES						
O		1.44	1.54	1.80	1.98	1.69
D		1.34	1.93	2.00	2.13	1.85
N*		0.89	1.28	1.44	1.39	1.25
PK		1.14	1.92	1.82	2.20	1.77
N*PK		0.81	1.66	1.76	1.64	1.47
Mean		1.12	1.67	1.76	1.87	1.60

STRAW MEAN DM% 90.2

PLOT AREA HARVESTED 0.00588

91/R/PG/5

PARK GRASS

Object: To study the effects of organic and inorganic manures and lime on old grass (for hay).

The 136th year, hay.

For previous years see 'Details' 1967 and 1973 and 74-90/R/PG/5.

Treatments: Combinations of:-

Whole plots

1. MANURE Fertilizers and organic manures:

N1	Plot 1	N1
O(D)	Plot 2	None (D until 1863)
O/PLOT3	Plot 3	None
P	Plot 4/1	P
N2P	Plot 4/2	N2 P
N1MN	Plot 6	N1 P K Na Mg
MN	Plot 7	P K Na Mg
PNAMG	Plot 8	P Na Mg
MN(N2)	Plot 9/1	P K Na Mg (N2 until 1989)
N2MN	Plot 9/2	N2 P K Na Mg
N2PNAMG	Plot 10	N2 P Na Mg
N3MN	Plot 11/1	N3 P K Na Mg
N3MNSI	Plot 11/2	N3 P K Na Mg Si
O/PLOT12	Plot 12	None
D/F	Plot 13	D/F
MN(N2*14)	Plot 14/1	P K Na Mg (N2* until 1989)
N2*MN	Plot 14/2	N2* P K Na Mg
MN(N2*15)	Plot 15	P K Na Mg (N2* until 1875)
N1*MN	Plot 16	N1* P K Na Mg
N1*	Plot 17	N1*
N2KNAMG	Plot 18	N2 K Na Mg
D	Plot 19	D
D/N*PK	Plot 20	D/N*P K
N1, N2, N3:	48, 96, 144 kg N as sulphate of ammonia	
N1*, N2*:	48, 96 kg N as nitrate of soda (30 kg N to Plot 20, only in years with no farmyard manure)	
P:	35 kg P (15 kg P to Plot 20, only in years with no farmyard manure) as single superphosphate until 1986, triple superphosphate in 1974, and since 1987	
K:	225 kg K (45 kg K to Plot 20, only in years with no farmyard manure) as sulphate of potash	
Na:	15 kg Na as sulphate of soda	
Mg:	10 kg Mg as sulphate of magnesia	
Si:	Silicate of soda at 450 kg	
D:	Farmyard manure at 35 tonnes every fourth year	
F:	Fish meal every fourth year to supply 63 kg N	
MN:	P K Na Mg	

91/R/PG/5

Sub plots

2. **LIME** Liming:

- | | |
|---|--|
| A | a Ground chalk applied as necessary to achieve pH7 |
| B | b Ground chalk applied as necessary to achieve pH6 |
| C | c Ground chalk applied as necessary to achieve pH5 |
| D | d None |

NOTE: Lime was applied regularly, and at the same rate, to all 'a' and 'b' sub plots of Plots 1 to 17 (except 12) from 1924. Differential liming started in 1965 on certain 'b' and 'c' sub plots (except on Plot 12) and in 1976 on certain 'a' sub plots (including Plot 12) and 12b. Liming ceased on plots 9/1 and 14/1 after 1989. Lime last applied in 1990.

Additional sub plots (Plots 18, 19 and 20 only) (tonnes CaCO₃ applied every fourth year 1920-1964):

N2KNAMG0	18-1	None
N2KNAMG2	18-2	13.5
N2KNAMG1	18-3	7.9
DO	19-1	None
D2	19-2	6.3
D1	19-3	1.1
D/N*PK0	20-1	None
D/N*PK2	20-2	5.6
D/N*PK1	20-3	1.1

Since 1965 Plot 18-1 has been split into two for treatments 'c' and 'd' above and Plot 18-3 split into two for treatments 'a' and 'b'. Plots 19 and 20 received no further chalk after 1968; plot 18/2 no further chalk after 1972.

NOTE: For a fuller record of treatments see 'Details' etc.

Cultivations, etc.:- P applied: 21 Nov, 1990. Fish meal applied: 28 Nov. K, Na, Mg and Si applied: 30 Nov. N applied: 10 Apr, 1991. Cut: 9 July, 6 Nov.

91/R/PG/5

1ST CUT (9/7/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

LIME MANURE	A	B	C	D	MEAN
N1	2.08	4.49	2.64	1.12	2.58
O(D)	3.31	4.27	2.51	2.82	3.23
O/PLOT3	3.48	3.78	2.53	2.74	3.13
P	4.30	4.20	4.00	4.05	4.14
N2P	4.65	4.09	7.37	4.00	5.03
N1MN	6.01	6.07			6.04
MN	5.56	5.40	4.93	5.09	5.25
PNAMG	3.25	3.83	4.79	4.79	4.17
MN(N2)	3.41	2.82	1.83	1.97	2.51
N2MN	5.93	6.05	6.09	6.26	6.08
N2PNAMG	4.42	4.37	4.58	3.56	4.23
N3MN	6.54	6.85	5.69	6.84	6.48
N3MNSI	6.80	6.35	6.28	6.43	6.47
O/PLOT12	2.48	3.12	2.46	2.90	2.74
D/F	5.60	5.58	5.59	4.96	5.43
MN(N2*14)	4.37	4.70	3.61	4.40	4.27
N2*MN	5.78	7.16	5.65	5.04	5.91
MN(N2*15)	6.19	5.32	4.42	4.19	5.03
N1*MN	5.99	5.26	4.82	4.70	5.19
N1*	4.06	4.07	4.41	4.07	4.15
N2KNAMG0			3.20	1.19	2.19
N2KNAMG2	4.52				4.52
N2KNAMG1	4.37	4.45			4.41
D0	5.22				5.22
D2	4.97				4.97
D1	5.46				5.46
D/N*PK0	5.40				5.40
D/N*PK2	5.45				5.45
D/N*PK1	5.40				5.40

1ST CUT MEAN DM% 25.5

91/R/PG/5

2ND CUT (6/11/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

LIME MANURE	A	B	C	D	MEAN
N1	2.15	2.41	1.26	0.65	1.62
O(D)	1.82	2.18	1.37	1.77	1.78
O/PLOT3	1.56	2.04	1.90	2.22	1.93
P	1.81	1.99	2.30	2.31	2.10
N2P	2.00	1.41	1.13	0.75	1.32
N1MN	2.61	2.81			2.71
MN	2.99	3.14	2.04	1.92	2.52
PNAMG	1.53	1.81	2.35	2.25	1.99
MN(N2)	1.53	1.11	0.57	0.82	1.01
N2MN	2.00	2.25	0.77	0.80	1.46
N2PNAMG	1.28	1.13	0.87	0.88	1.04
N3MN	2.77	1.66	0.87	2.91	2.05
N3MNSI	2.65	2.33	1.65	2.69	2.33
O/PLOT12	0.92	1.39	1.26	1.60	1.29
D/F	2.76	2.89	2.64	2.81	2.77
MN(N2*14)	2.25	1.93	1.53	1.75	1.87
N2*MN	2.44	2.40	2.21	1.96	2.25
MN(N2*15)	2.51	2.54	2.65	2.13	2.46
N1*MN	2.58	2.50	2.40	1.99	2.37
N1*	2.37	2.33	2.32	2.12	2.29
N2KNAMG0			1.58	0.43	1.00
N2KNAMG2	2.69				2.69
N2KNAMG1	3.18	2.57			2.87
D0	3.34				3.34
D2	3.34				3.34
D1	3.04				3.04
D/N*PK0	3.06				3.06
D/N*PK2	2.83				2.83
D/N*PK1	3.73				3.73

2ND CUT MEAN DM% 25.2

91/R/PG/5

TOTAL OF 2 CUTS DRY MATTER TONNES/HECTARE

***** Tables of means *****

LIME MANURE	A	B	C	D	MEAN
N1	4.24	6.90	3.90	1.76	4.20
O(D)	5.13	6.45	3.88	4.59	5.01
O/PLOT3	5.04	5.83	4.44	4.95	5.06
P	6.11	6.19	6.30	6.36	6.24
N2P	6.65	5.51	8.49	4.75	6.35
N1MN	8.62	8.88			8.75
MN	8.56	8.54	6.97	7.01	7.77
PNAMG	4.78	5.64	7.14	7.05	6.15
MN(N2)	4.94	3.93	2.40	2.79	3.52
N2MN	7.93	8.30	6.86	7.06	7.54
N2PNAMG	5.70	5.50	5.45	4.44	5.27
N3MN	9.31	8.52	6.57	9.74	8.54
N3MNSI	9.44	8.68	7.93	9.12	8.80
O/PLOT12	3.39	4.51	3.72	4.50	4.03
D/F	8.36	8.47	8.23	7.77	8.21
MN(N2*14)	6.62	6.63	5.14	6.15	6.14
N2*MN	8.22	9.56	7.86	7.00	8.16
MN(N2*15)	8.70	7.86	7.07	6.32	7.49
N1*MN	8.56	7.75	7.22	6.69	7.56
N1*	6.43	6.40	6.73	6.19	6.44
N2KNAMG0			4.78	1.62	3.20
N2KNAMG2	7.21				7.21
N2KNAMG1	7.55	7.02			7.28
D0	8.56				8.56
D2	8.31				8.31
D1	8.50				8.50
D/N*PK0	8.46				8.46
D/N*PK2	8.28				8.28
D/N*PK1	9.14				9.14

TOTAL OF 2 CUTS MEAN DM% 25.4

PLOT AREA HARVESTED 0.00002

91/R/BN/7

BARNFIELD

Object: The experiment was designed to study the effects of organic and inorganic manures on continuous root crops. It was progressively modified to study effects on other crops.

Sections 1 and 2 the eighth year of grass/clover. The 17th year of grass on the rest of the experiment.

For previous years see 'Details' 1967 and 1973 and 74-90/R/BN/7.

Plot dimensions: 10.7 x 55.9.

Treatments to grass: All combinations of:-

Whole plots

1. **MANURE** Fertilizers and organic manures:

D	D
DPK	D P K
PKMG	P K (Na) Mg
P	P
PK	P K
PMG	P (Na) Mg
0	0

- P: 35 kg P as single superphosphate until 1987, triple superphosphate since and in 1974
K: 225 kg K as sulphate of potash
(Na): 90 kg Na as sodium chloride until 1973
Mg: 90 kg Mg as kieserite every fourth year since 1974 (sulphate of magnesia until 1973)
D: Farmyard manure at 35 tonnes (until 1975).

Quarter plots

2. **N PERCUT** Nitrogen fertilizer in 1991 (kg N per cut) as 'Nitram', cumulative to previous dressings, and residues of forms of N previously each supplying 96 kg N per annum:

75	75, previously nitrate of soda, section 3
100	100, previously sulphate of ammonia, section 4
125	125, previously sulphate of ammonia + castor meal, section 5
150	150, previously castor meal, section 6

Castor meal last applied 1961, nitrate of soda and sulphate of ammonia until 1959.

Plus one plot **MANURE** KMG 100

91/R/BN/7

Treatments to grass/clover, sections 1 and 2 (not given nitrogen fertilizer):

MANURE Fertilizers and organic manures as for grass above, excluding KMG.

NOTES: (1) P, K and D treatments were applied to Sections 1 and 2 until 1980. None were applied subsequently until the resumption of P and K treatments, only, from 1985.
(2) Yields were not taken from section 2.

Cultivations, etc.:-

All sections: P applied: 21 Nov, 1990. K applied: 28 Nov. Cut: 30 May, 1991 and 11 Nov.

Grass (Sections 3, 4, 5 and 6) only: N applied: 8 Mar, 1991 and 10 June.

GRASS

1ST CUT (30/5/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

N PERCUT MANURE	75	100	125	150	Mean
D	4.90	4.58	4.63	4.72	4.71
DPK	4.63	4.65	5.07	4.98	4.83
PKMG	4.47	4.64	4.50	4.72	4.58
P	2.25	1.81	1.65	1.69	1.85
PK	4.72	4.15	4.57	4.62	4.51
PMG	2.50	1.96	1.75	1.80	2.00
0	2.55	2.36	1.94	2.06	2.22
Mean	3.72	3.45	3.45	3.51	3.53

MANURE KMG 100 3.73

Grand mean 3.54

1ST CUT MEAN DM% 21.6

91/R/BN/7

GRASS

2ND CUT (11/11/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

N PERCUT MANURE	75	100	125	150	Mean
D	5.15	5.88	4.78	5.60	5.35
DPK	6.12	5.80	5.66	5.39	5.74
PKMG	5.24	4.04	4.73	4.85	4.72
P	2.94	1.52	1.76	1.81	2.01
PK	4.91	5.12	5.03	5.07	5.03
PMG	2.65	1.53	1.17	1.49	1.71
0	2.56	2.51	2.41	2.48	2.49
Mean	4.22	3.77	3.65	3.81	3.86

MANURE KMG 100 4.63

Grand mean 3.89

2ND CUT MEAN DM% 27.2

TOTAL OF 2 CUTS DRY MATTER TONNES/HECTARE

***** Tables of means *****

N PERCUT MANURE	75	100	125	150	Mean
D	10.05	10.46	9.42	10.33	10.06
DPK	10.75	10.45	10.73	10.37	10.57
PKMG	9.71	8.68	9.23	9.56	9.30
P	5.19	3.32	3.41	3.50	3.86
PK	9.63	9.28	9.60	9.68	9.55
PMG	5.15	3.50	2.93	3.29	3.72
0	5.11	4.86	4.34	4.53	4.71
Mean	7.94	7.22	7.09	7.33	7.39

MANURE KMG 100 8.36

Grand mean 7.43

TOTAL OF 2 CUTS MEAN DM% 24.4

PLOT AREA HARVESTED 0.00568

91/R/BN/7

GRASS/CLOVER

1ST CUT (30/5/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

MANURE	D	DPK	PKMG	P	PK	PMG	0	Mean
	1.18	1.21	0.57	0.25	0.34	0.28	0.29	0.59

1ST CUT MEAN DM% 30.3

2ND CUT (11/11/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

MANURE	D	DPK	PKMG	P	PK	PMG	0	Mean
	1.59	1.62	1.91	1.03	2.21	1.70	1.47	1.65

2ND CUT MEAN DM% 22.7

TOTAL OF 2 CUTS DRY MATTER TONNES/HECTARE

***** Tables of means *****

MANURE	D	DPK	PKMG	P	PK	PMG	0	Mean
	2.77	2.83	2.48	1.28	2.55	1.99	1.76	2.24

TOTAL OF 2 CUTS MEAN DM% 26.5

PLOT AREA HARVESTED 0.00568

91/R/GC/8

GARDEN CLOVER

Object: To study yields and pathogens of red clover grown continuously - Manor Garden.

The 138th year, red clover.

For previous years see 'Details' 1967 and 1973, and 74-90/R/GC/8.

Design: 2 blocks of 2 plots.

Whole plot dimensions: 1.00 x 1.40.

Treatments:

FUNG RES Residual effects of fungicide to control *Sclerotinia trifoliorum*:

NONE None

BENOMYL Benomyl sprays during previous winters, last applied November 1989.

Basal applications: Manures: Chalk at 1.25 t. (0:18:36) at 420 kg.
Mg at 50 kg, as Epsom Salts.

NOTE: Additional K was applied to replace that removed by the crop in 1990. **FUNG RES** NONE required 157 and 129 kg K2O to the first and second blocks respectively, **FUNG RES BENOMYL** 136 and 124 kg K2O. This was applied as muriate of potash, one third in spring 1991 and one third after the first and second cuts.

Seed: Hungaropoly, sown at 30 kg in 1990.

Cultivations, etc.:- Chalk, P, K and Mg applied: 23 Oct, 1990. Hand weeded: 13 Mar, 1991. K applied: 11 Apr. Cut, weeded and K applied: 5 June, 25 July. Cut and weeded: 20 Sept.

91/R/GC/8

1ST CUT (5/6/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

FUNG RES	NONE	BENOMYL	Mean
	8.31	7.86	8.09

1ST CUT MEAN DM% 14.5

2ND CUT (25/7/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

FUNG RES	NONE	BENOMYL	Mean
	6.98	6.12	6.55

2ND CUT MEAN DM% 13.5

3RD CUT (20/9/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

FUNG RES	NONE	BENOMYL	Mean
	5.06	4.11	4.58

3RD CUT MEAN DM% 18.7

TOTAL OF 3 CUTS DRY MATTER TONNES/HECTARE

***** Tables of means *****

FUNG RES	NONE	BENOMYL	Mean
	20.35	18.09	19.22

TOTAL OF 3 CUTS MEAN DM% 15.6

PLOT AREA HARVESTED 0.00010