


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1991

[Full Table of Content](#)



Sunflowers

Rothamsted Research

Rothamsted Research (1992) *Sunflowers* ; Yields Of The Field Experiments 1991, pp 159 - 162 -
DOI: <https://doi.org/10.23637/ERADOC-1-46>

91/R/SU/1

SUNFLOWERS

VARIETIES AND BOTRYTIS

Object: To study the incidence of *Botrytis cinerea* on two varieties of sunflower and to compare their yields in the absence of fungicide treatment - Great Harpenden I.

Sponsors: V.J. Church, B.D.L. Fitt.

Design: 8 randomised blocks of 2 plots.

Whole plot dimensions: 3.5 x 10.0.

Treatments:

VARIETY Varieties:

S 47

ALLEGRO

Basal applications: Manures: (0:16:36) at 980 kg. (12:20:20) at 380 kg.
Weedkillers: Trifluralin at 1.1 kg in 200 l. Linuron at 0.48 kg in 200 l.

Seed: Varieties, sown at 12 seeds per square metre.

Cultivations, etc.:- First PK applied: 3 Sept, 1990. Ploughed: 11 Dec.
NPK applied: 11 Apr, 1991. Heavy spring-tine cultivated: 24 Apr.
Rotary harrowed, trifluralin applied, rotary harrowed, seed sown, rolled: 29 Apr. Linuron applied: 9 May. Hand harvested: 9 Sept.
Previous crops: W. barley 1989 and 1990.

NOTE: Selected plants were monitored daily for growth stage and for disease from flowering to harvest. Numbers of *Botrytis* spores were recorded daily using a Burkard spore trap. Seed moisture content was measured at intervals after flowering.

91/R/SU/1

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

VARIETY

S 47 2.48

ALLEGRO 2.43

Mean 2.45

*** Standard errors of differences of means ***

VARIETY

0.071

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
---------	------	------	-----

BLOCK.WP	7	0.143	5.8
----------	---	-------	-----

GRAIN MEAN DM% 71.2

PLOT AREA HARVESTED 0.00150

91/R/SU/2

SUNFLOWERS

BIRD REPELLANT

Object: To study the effects of applying two rates of a saponin, at two times, on damage by birds to and yields of sunflowers - Great Harpenden I.

Sponsor: V.J. Church.

Design: 5 x 5 quasi-complete Latin square.

Whole plot dimensions: 3.5 x 10.0.

Treatments:

SAP R T Rates and times of applying a saponin (as weight of quinoa husk):

NONE	None
S1 E	51 kg applied 8 Aug, 1991
S1 L	51 kg applied 28 Aug
S2 E	57 kg applied 8 Aug
S2 L	57 kg applied 28 Aug

NOTES: (1) On S1 plots saponin was applied as ground quinoa husk in 860 l/ha.
(2) On S2 plots saponin was applied as ground quinoa husk as a dry dust.

Basal applications: Manures: (0:16:36) at 980 kg. (12:20:20) at 380 kg.
Weedkillers: Trifluralin at 1.1 kg in 200 l. Linuron at 0.50 kg, in 400 l.

Seed: Allegro, sown at 9 seeds per square metre.

Cultivations, etc.:- First PK applied: 3 Sept, 1990. Ploughed: 11 Dec.
NPK applied: 11 Apr, 1991. Heavy spring-tine cultivated: 24 Apr.
Rotary harrowed, trifluralin applied, rotary harrowed, seed sown, rolled: 26 Apr. Linuron applied: 3 May. Hand harvested: 10 Sept..
Previous crops: W. barley 1989 and 1990.

NOTE: Growth stage and bird damage was monitored twice weekly. Rainfall was recorded after repellent application. Disease was assessed before harvest and seed moisture measured on three occasions before harvest.

91/R/SU/2

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

SAP R T	
NONE	1.91
S1 E	2.08
S1 L	1.84
S2 E	1.80
S2 L	2.03
Mean	1.93

*** Standard errors of differences of means ***

SAP R T
0.115

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
ROW.COL	12	0.182	9.4
GRAIN MEAN DM%	62.4		
PLOT AREA HARVESTED	0.00150		