


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1992

[Full Table of Content](#)



92/R/EX/4 Exhaustion Land - W. Wheat

Rothamsted Research

Rothamsted Research (1993) *92/R/EX/4 Exhaustion Land - W. Wheat* ; Yields Of The Field Experiments 1992, pp 22 - 24 - DOI: <https://doi.org/10.23637/ERADOC-1-47>

92/R/EX/4

EXHAUSTION LAND

Object: To study the residual effects of manures applied 1876-1901, and of additional phosphate applied since 1986, on the yield of continuous s. barley up to 1991, w. wheat in 1992 - Hoosfield.

The 137th year, w. wheat.

For previous years see 'Details' 1967, 1973 and 74-91/R/EX/4.

Treatments: All combinations of:-

Whole plots (P test)

1. **OLD RES** Residues of manures applied annually 1876-1901:

O	None
D	Farmyard manure at 35 tonnes
N	96 kg N as ammonium salts
P	34 kg P as superphosphate
NPKNAMG	N and P as above plus 137 kg K as sulphate of potash, 16 kg Na as sulphate of soda, 11 kg Mg as sulphate of magnesia

2. **P** Phosphate applied annually from 1986 as
 superphosphate until 1987, triple superphosphate
 since:

O	None
P1	44 kg P
P2	87 kg P
P3	131 kg P

plus

Whole plots (K test, previously N test until 1991)

- | | |
|----------------|--|
| OLD RES | Residues of manures applied annually 1876-1901: |
| O | None |
| D | Farmyard manure at 35 tonnes |
| N* | 96 kg N as nitrate of soda |
| PK | 34 kg P as superphosphate, 137 kg K as sulphate of
potash |
| N*PK | N, P and K as above |

Experimental diary:

P test:

18-Sep-91 : **T** : P applied as triple superphosphate to treatment plots.
 : **T** : Muriate of potash at 170 kg.

Residual N test:

18-Sep-91 : **T** : Triple superphosphate at 1420 kg.
23-Sep-91 : **T** : Triple superphosphate at 710 kg.

92/R/EX/4

Experimental diary:

All plots:

19-Sep-91 : B : Sting CT at 1.5 l in 200 l.
 20-Sep-91 : B : Ploughed, spring-tine cultivated, rolled.
 23-Sep-91 : B : Rotary harrowed, Mercia drilled at 160 kg, rolled.
 14-Apr-92 : B : 34.5% N applied at 560 kg.
 23-Jun-92 : B : Radar at 0.50 l and Mistral at 0.50 l in 200 l.
 31-Jul-92 : B : Combine harvested.

NOTES: (1) Yields presented for **K TEST** are means of four plots.
 (2) Grain and straw samples were taken for chemical analysis.

P TEST

GRAIN TONNES/HECTARE

***** Tables of means *****

	P	O	P1	P2	P3	Mean
OLD RES						
O		2.70	7.97	8.17	7.96	6.70
D		6.85	8.21	8.03	8.11	7.80
N		2.43	7.94	7.92	8.11	6.60
P		5.08	7.46	7.26	7.63	6.86
NPKNAMG		4.69	7.70	7.90	7.84	7.03
Mean		4.35	7.86	7.86	7.93	7.00

GRAIN MEAN DM% 87.5

STRAW TONNES/HECTARE

***** Tables of means *****

	P	O	P1	P2	P3	Mean
OLD RES						
O		1.71	6.39	6.96	6.79	5.46
D		4.72	6.60	7.21	6.43	6.24
N		1.87	6.62	7.50	7.18	5.79
P		4.27	6.03	8.13	7.60	6.51
NPKNAMG		3.82	6.39	7.20	6.91	6.08
Mean		3.28	6.41	7.40	6.98	6.02

STRAW MEAN DM% 85.1

PLOT AREA HARVESTED 0.00589

92/R/EX/4

K TEST

GRAIN TONNES/HECTARE

***** Tables of means *****

OLD RES	O	D	N*	PK	N*PK	Mean
	5.70	7.35	5.25	4.96	5.69	5.79

GRAIN MEAN DM% 87.8

STRAW TONNES/HECTARE

***** Tables of means *****

OLD RES	O	D	N*	PK	N*PK	Mean
	5.24	5.70	4.90	5.55	5.19	5.32

STRAW MEAN DM% 91.6

PLOT AREA HARVESTED 0.00589