

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1994

[Full Table of Content](#)



94/R/CS/10 and 94/W/CS/10 Long-term Liming - W. Lupins

Rothamsted Research

Rothamsted Research (1995) *94/R/CS/10 and 94/W/CS/10 Long-term Liming - W. Lupins* ; Yields Of The Field Experiments 1994, pp 52 - 53 - DOI: <https://doi.org/10.23637/ERADOC-1-49>

94/R/CS/10 and 94/W/CS/10

LONG TERM LIMING

Object: To study the effects of different amounts of lime, phosphate and sulphur on the yields and compositions of a sequence of crops - Rothamsted (R) Sawyers I and Woburn (W) Stackyard C.

Sponsors: S.P. McGrath, P.B. Barraclough, G.F.J. Milford, J.M. Day.

The 33rd year, w. lupins.

For previous years see 'Details' 1967, 1973 and 74-93/R&W/CS/10.

Design: 2 randomised blocks of 16 plots split into 2 sub-plots.

Whole plot dimensions: 5.8 x 16.1 (R), 5.6 x 16.1 (W).

Treatments: All combinations of:-

Whole plots

1. **CHALK** Residual effects of ground chalk (tonnes CaCO₃) (total applied 1962-87):

R	W	Rothamsted total		Woburn total	
		1962-78	1982-87	1962-78	1982-87
0	0	0	0	0	0
15	9	7	8	6	3
24.5	25.5	15	9.5	14	11.5
52.5	45.5	30	22.5	23	22.5

2. **P** Residual effects of P fertilizer applied:

	Until 1978	1981	1982	1983		1988	
	R & W	R & W	R & W	R	W	R	W
0	0	0	0	0	0	0	0
P1	0	P1	P1	0	P2	P1	P1
P2	P	P1	0	P2	P2	P1	P1
P3	P	P3	P1	P2	P4	P3	P3

Rates 1981-83 and 1988 P1, P2, P3, P4 = 25, 50, 75, 100 kg P as superphosphate

Sub-plots

3. **SULPHUR** Sulphur (kg S, as calcium sulphate), applied cumulatively since 1991:

0
30

94/R/CS/10 and 94/W/CS/10

NOTES: (1) Until 1978 test P was applied cumulatively, rates varied with crop, none in 1979 and 1980. K was also applied cumulatively, to P1 and P3 plots. Since 1981 K has been applied basally (none in 1986, 1987, 1989, 1990, 1993 and 1994).

(2) Test manganese was applied cumulatively, 1987-90.

Experimental diary:

Sawyers I (R):

16-Oct-93 : B : Ploughed.

19-Oct-93 : B : Discd.

20-Oct-93 : B : Spring-tine cultivated, CH304/70 drilled at 40 seeds per m².

26-Oct-93 : B : Opogard 500 FW at 2.8 l in 200 l.

16-Feb-94 : B : Atlas Simazine at 1.0 l in 200 l.

13-Jun-94 : T : **SULPHUR** 30: 30 kg S as gypsum.

06-Sep-94 : B : Barclay Gallup at 4.0 l in 200 l.

Stackyard C (W):

16-Sep-93 : B : Ploughed.

24-Sep-93 : B : Rotary harrowed, CH304/70 drilled at 40 seeds per m².

18-Feb-94 : B : Atlas Simazine at 1.0 l in 200 l.

10-Mar-94 : T : **SULPHUR** 30: 30 kg S as gypsum.

09-May-94 : B : Sportak 45 at 1.5 l in 200 l.

NOTE: Due to poor winter survival at Rothamsted and to excessive grazing damage at Woburn, both crops were abandoned in summer and no yields were taken.