

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1994

[Full Table of Content](#)



94/R/CS/140 Chemical Reference Plots - S. Barley

Rothamsted Research

Rothamsted Research (1995) *94/R/CS/140 Chemical Reference Plots - S. Barley* ; Yields Of The Field Experiments 1994, pp 54 - 57 - DOI: <https://doi.org/10.23637/ERADOC-1-49>

94/R/CS/140

CHEMICAL REFERENCE PLOTS

Object: To study the persistence in soil of agricultural chemicals applied annually, singly and in combination, and their effects on soil microflora and on yield of continuous s. barley - Long Hoos V 3.

Sponsors: R.H. Bromilow, A.A. Evans, P.H. Nicholls.

The 21st year, s. barley.

For previous years see 74-93/R/CS/140.

Design: Single replicate of 32 plots.

Whole plot dimensions: 4.06 x 4.57.

Treatments: Applied cumulatively every year until 1993, none in 1994.

All combinations of:-

1. **WEEDKLLR** Weedkiller in autumn:
(NONE) None
(GLYPHOS) Glyphosate to barley stubble each autumn
2. **FUNGICIDE[1]** Fungicide in autumn:
(NONE) None
(TRIADIM) Triadimefon in autumn
3. **FUNGICIDE[2]** Fungicide in spring:
(NONE) None
(BENOMYL) Benomyl to the seedbed
4. **INSECTCDE** Insecticide:
(NONE) None
(CHLORFEN) Chlorfenvinphos to the seedbed
5. **NEMACIDE** Nematicide:
(NONE) None
(ALDICARB) Aldicarb to the seedbed

Experimental diary:

03-Nov-93 : B : Ploughed.
17-Mar-94 : B : Spring-tine cultivated, rotary harrowed, Alexis
undressed, drilled at 350 seeds per m², rolled.
19-Apr-94 : B : 34.5% N at 428 kg.
02-Jun-94 : B : Oxytril CM at 1.5 l with Duplosan New System CMPP at
2.0 l in 200 l.
08-Aug-94 : B : Combine harvested.

94/R/CS/140

NOTE: Samples of topsoil were taken from representative plots in August 1994 for the estimation of pesticide residues. The influence of treatment history on the breakdown rates of the chemicals was examined in these samples.

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNGCIDE[1]	(NONE)	(TRIADIM)	Mean
WEEDKLLR			
(NONE)	5.29	5.35	5.32
(GLYPHOS)	5.19	5.25	5.22
Mean	5.24	5.30	5.27

FUNGCIDE[2]	(NONE)	(BENOMYL)	Mean
WEEDKLLR			
(NONE)	5.25	5.39	5.32
(GLYPHOS)	5.08	5.36	5.22
Mean	5.16	5.37	5.27

FUNGCIDE[2]	(NONE)	(BENOMYL)	Mean
FUNGCIDE[1]			
(NONE)	5.24	5.24	5.24
(TRIADIM)	5.09	5.51	5.30
Mean	5.16	5.37	5.27

INSTCDE	(NONE)	(CHLORFEN)	Mean
WEEDKLLR			
(NONE)	5.45	5.20	5.32
(GLYPHOS)	5.34	5.09	5.22
Mean	5.39	5.14	5.27

INSTCDE	(NONE)	(CHLORFEN)	Mean
FUNGCIDE[1]			
(NONE)	5.24	5.23	5.24
(TRIADIM)	5.55	5.05	5.30
Mean	5.39	5.14	5.27

INSTCDE	(NONE)	(CHLORFEN)	Mean
FUNGCIDE[2]			
(NONE)	5.24	5.09	5.16
(BENOMYL)	5.55	5.20	5.37
Mean	5.39	5.14	5.27

94/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

NEMACIDE	(NONE)	(ALDICARB)	Mean
WEEDKLLR			
(NONE)	5.39	5.26	5.32
(GLYPHOS)	5.25	5.18	5.22
Mean	5.32	5.22	5.27
NEMACIDE	(NONE)	(ALDICARB)	Mean
FUNGICIDE[1]			
(NONE)	5.31	5.17	5.24
(TRIADIM)	5.32	5.28	5.30
Mean	5.32	5.22	5.27
NEMACIDE	(NONE)	(ALDICARB)	Mean
FUNGICIDE[2]			
(NONE)	5.20	5.13	5.16
(BENOMYL)	5.43	5.31	5.37
Mean	5.32	5.22	5.27
NEMACIDE	(NONE)	(ALDICARB)	Mean
INSCTCDE			
(NONE)	5.39	5.40	5.39
(CHLORFEN)	5.24	5.05	5.14
Mean	5.32	5.22	5.27
FUNGICIDE[1]	(NONE)	(TRIADIM)	
WEEDKLLR FUNGICIDE[2]	(NONE)	(BENOMYL)	(NONE) (BENOMYL)
(NONE)	5.23	5.35	5.27 5.43
(GLYPHOS)	5.25	5.12	4.91 5.59
FUNGICIDE[1]	(NONE)	(TRIADIM)	
WEEDKLLR INSCTCDE	(NONE)	(CS)	(NONE) (CS)
(NONE)	5.35	5.23	5.54 5.16
(GLYPHOS)	5.14	5.23	5.55 4.94
FUNGICIDE[2]	(NONE)	(BENOMYL)	
WEEDKLLR INSCTCDE	(NONE)	(CS)	(NONE) (CS)
(NONE)	5.23	5.27	5.66 5.13
(GLYPHOS)	5.25	4.90	5.43 5.28
FUNGICIDE[2]	(NONE)	(BENOMYL)	
FUNGICIDE[1] INSCTCDE	(NONE)	(CS)	(NONE) (CS)
(NONE)	5.13	5.35	5.36 5.12
(TRIADIM)	5.36	4.82	5.73 5.29

94/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

	FUNGCIDE[1]	(NONE)		(TRIADIM)	
WEEDKLLR	NEMACIDE	(NONE)	(AL)	(NONE)	(AL)
(NONE)		5.38	5.20	5.39	5.31
(GLYPHOS)		5.24	5.13	5.25	5.24

	FUNGCIDE[2]	(NONE)		(BENOMYL)	
WEEDKLLR	NEMACIDE	(NONE)	(AL)	(NONE)	(AL)
(NONE)		5.30	5.20	5.47	5.31
(GLYPHOS)		5.10	5.05	5.39	5.32

	FUNGCIDE[2]	(NONE)		(BENOMYL)	
FUNGCIDE[1]	NEMACIDE	(NONE)	(AL)	(NONE)	(AL)
(NONE)		5.31	5.17	5.31	5.16
(TRIADIM)		5.09	5.09	5.55	5.47

	INSCTCDE	(NONE)		(CHLORFEN)	
WEEDKLLR	NEMACIDE	(NONE)	(AL)	(NONE)	(AL)
(NONE)		5.46	5.43	5.32	5.08
(GLYPHOS)		5.33	5.36	5.17	5.01

	INSCTCDE	(NONE)		(CHLORFEN)	
FUNGCIDE[1]	NEMACIDE	(NONE)	(AL)	(NONE)	(AL)
(NONE)		5.27	5.21	5.35	5.12
(TRIADIM)		5.51	5.58	5.14	4.97

	INSCTCDE	(NONE)		(CHLORFEN)	
FUNGCIDE[2]	NEMACIDE	(NONE)	(AL)	(NONE)	(AL)
(NONE)		5.20	5.28	5.20	4.97
(BENOMYL)		5.58	5.51	5.28	5.12

*** Standard errors of differences of means ***

Margins of two factor tables 0.109
 Two factor tables 0.154
 Three factor tables 0.218

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
WP	6	0.308	5.8

GRAIN MEAN DM% 87.9

PLOT AREA HARVESTED 0.00069