


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1994

[Full Table of Content](#)



94/R/WW/1 Nitrogen Indicators - W. Wheat

Rothamsted Research

Rothamsted Research (1995) *94/R/WW/1 Nitrogen Indicators - W. Wheat* ; Yields Of The Field Experiments 1994, pp 93 - 94 - DOI: <https://doi.org/10.23637/ERADOC-1-49>

94/R/WW/1

WINTER WHEAT

NITROGEN INDICATORS

Object: To relate chlorophyll concentrations in individual leaves of w. wheat to nitrogen supply and crop yield - Delharding.

Sponsor: P.B. Barraclough.

Design: 3 randomised blocks of 8 plots.

Plot dimensions: 3.0 x 15.0.

Treatments:

N	Nitrogen in spring (kg N) applied as 34.5% N at first node formation:
-	0
1	50
2	100
3	150
4	200
5	250
6	300
7	40 and, in addition, four subsequent dressings of 40, applied at weekly intervals (total 200)

Experimental diary:

24-Aug-93 : B : Ploughed and furrow pressed.
27-Aug-93 : B : Rolled.
15-Oct-93 : B : Triple superphosphate at 213 kg.
18-Oct-93 : B : Spring-tine cultivated.
23-Oct-93 : B : Rotary harrowed, Hereward, dressed Cerevax, drilled at 380 seeds per m².
18-Apr-94 : T : N 1: 34.5% N at 145 kg.
 : T : N 2: 34.5% N at 290 kg.
 : T : N 3: 34.5% N at 435 kg.
 : T : N 4: 34.5% N at 580 kg.
 : T : N 5: 34.5% N at 725 kg.
 : T : N 6: 34.5% N at 870 kg.
 : T : N 7: 34.5% N at 116 kg.
25-Apr-94 : T : N 7: 34.5% N at 116 kg.
01-May-94 : B : Halo at 2.0 l with Tripart Brevis at 2.5 l in 200 l.
03-May-94 : T : N 7: 34.5% N at 116 kg.
09-May-94 : B : Ally at 30 g with Cheetah Super at 3.0 l and Starane 2 at 0.75 l in 200 l.
10-May-94 : T : N 7: 34.5% N at 116 kg.
18-May-94 : T : N 7: 34.5% N at 116 kg.
23-Jun-94 : B : Cyclone at 1.0 l with Mallard 750 EC at 0.5 l in 200 l.
22-Aug-94 : B : Combine harvested.

94/R/WW/1

- NOTES: (1) Leaf chlorophyll was measured at weekly intervals with a hand-held meter.
 (2) Crop growth, leaf area index and N content of the crop were measured on four occasions in spring and summer.
 (3) Yield components and N content of straw and grain were measured at final harvest.

GRAIN TONNES/HECTARE

***** Tables of means *****

N	
-	5.28
1	7.22
2	8.88
3	9.29
4	9.36
5	10.05
6	9.90
7	9.93
Mean	8.74

*** Standard errors of differences of means ***

N
0.432

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	14	0.529	6.0
GRAIN MEAN DM%	87.9		
PLOT AREA HARVESTED	0.00253		