


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1995

[Full Table of Content](#)



95/R/WW/1 Prediction of Weed Competition - W. Wheat

Rothamsted Research

Rothamsted Research (1996) *95/R/WW/1 Prediction of Weed Competition - W. Wheat* ; Yields Of The Field Experiments 1995, pp 97 - 98 - DOI: <https://doi.org/10.23637/ERADOC-1-50>

95/R/WW/1

WINTER WHEAT

PREDICTION OF WEED COMPETITION

Object: To predict the yield response of winter wheat to competition from three contrasting weed species - Stackyard.

Sponsors: J.W. Cussans, P.J.W. Lutman.

Design: 3 randomised blocks of 3 x 6 plots.

Whole plot dimensions: 3.0 x 8.0.

Treatments: All combinations of:-

1. WEED SP	Weed species:		
SM	<i>Stellaria media</i> (chickweed)		
AM	<i>Alopecurus myosuroides</i> (black-grass)		
GA	<i>Galium aparine</i> (cleavers)		
2. WEED DEN	Weed density, plants per m ² :		
	SM	AM	GA
0	0	0	0
2	65	130	9
4	80	233	8
8	151	264	22
16	245	434	29
32	301	607	47

NOTE: Target weed densities, plants per m²: SM, AM: 0, 40, 80, 160, 320 and 640, GA: 0, 3, 6, 12, 24 and 48.

Experimental diary:

- 02-Sep-94 : B : PK as (0:20:32) at 1317 kg.
- 07-Sep-94 : B : Ploughed and furrow pressed.
- 30-Sep-94 : B : Rotary harrowed.
- : T : Weed seeds sown by hand.
- : B : Rotary harrowed, Mercia, dressed Rappor, drilled at 275 seeds per m².
- 21-Nov-94 : B : Draza at 5.5 kg.
- 13-Jan-95 : T : **WEED SP** SM, GA (except **WEED DEN** 0): Cheetah R at 2.5 l in 220 l.
- : T : **WEED DEN** 0: Stefes IPU at 3.0 l with Stomp 400 at 3.3 l in 220 l.
- : T : **WEED SP** AM (except **WEED DEN** 0): Oxytril CM at 1.5 l with Starane 2 at 0.75 l in 220 l.
- 14-Mar-95 : B : 34.5% N at 118 kg.
- 21-Mar-95 : T : **WEED SP** GA (except **WEED DEN** 0): Isoproturon 500 at 3.0 l in 220 l.
- : T : **WEED DEN** 0: Ally at 30 g with Cheetah R at 2.5 l and Starane 2 at 0.75 l in 220 l.
- 19-Apr-95 : B : 34.5% N at 463 kg.

95/R/WW/1

Experimental diary:

05-May-95 : B : Calixin at 0.35 l with Halo at 2.0 l in 200 l.
 20-Jun-95 : B : Silvacur at 1.0 l in 200 l.
 04-Aug-95 : B : Hand harvested.

Previous crops: S. rape 1993, w. oats 1994.

NOTE: Weeds were counted in autumn and spring. Weeds and crop were sampled monthly for dry weight, green area and height, crop also had tillers and leaves per plant counted. Soil was sampled for nitrogen content in February. Flag leaf area and nutrient content were assessed in June. Black-grass seed produced was measured in June. Components of yield were assessed after harvest.

GRAIN TONNES/HECTARE

***** Tables of means *****

WEED DEN	0	2	4	8	16	32	Mean
WEED SP							
SM	7.92	5.97	6.62	5.63	4.59	5.32	6.01
AM	8.37	3.48	2.90	3.20	2.49	2.65	3.85
GA	8.34	7.17	6.95	7.21	5.87	4.96	6.75
Mean	8.21	5.54	5.49	5.35	4.31	4.31	5.53

*** Standard errors of differences of means ***

WEED SP	WEED DEN	WEED SP
		WEED DEN
0.221	0.313	0.543

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	34	0.664	12.0

GRAIN MEAN DM% 91.2

PLOT AREA HARVESTED 0.00020