

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1996

[Full Table of Content](#)



96/R/WW/6 Herbicide Resistant Black-grass - W. Wheat

Rothamsted Research

Rothamsted Research (1997) *96/R/WW/6 Herbicide Resistant Black-grass - W. Wheat* ; Yields Of The Field Experiments 1996, pp 112 - 114 - DOI: <https://doi.org/10.23637/ERADOC-1-51>

96/R/WW/6

WINTER WHEAT

HERBICIDE RESISTANT BLACK-GRASS

Object: To evaluate the efficacy of different herbicides on a herbicide-resistant black-grass population and to determine any changes in the degree of resistance in the progeny of surviving plants - Claycroft.

Sponsor: S.R. Moss.

Design: 4 blocks of 15 plots.

Whole plot dimensions: 3.0 x 12.0.

Treatments:

HERBICIDE	Herbicide type, rate of active ingredient and timing (black-grass growth stage):
1	None (duplicated)
3	Isoproturon at 2.5 kg at two leaf stage
4	Tri-allate at 2.24 kg pre-emergence, plus isoproturon at 2.5 kg at two leaf stage
5	Trifluralin at 0.96 kg pre-emergence, plus isoproturon at 2.5 kg at two leaf stage
6	Pendimethalin at 2.0 kg pre-emergence, plus isoproturon at 2.5 kg at two leaf stage
7	Tri-allate at 2.24 kg plus trifluralin at 0.96 kg pre- emergence, plus isoproturon at 2.5 kg at two leaf stage
8	Tri-allate at 2.24 kg plus trifluralin at 0.96 kg pre- emergence, isoproturon at 2.5 kg at two leaf stage and at 2.1 kg at one tiller stage
9	Tri-allate at 2.24 kg plus trifluralin at 0.96 kg pre- emergence, isoproturon at 2.5 kg at two leaf stage and fenoxaprop-P-ethyl at 0.069 kg at one tiller stage
10	Tri-allate at 2.24 kg plus trifluralin at 0.96 kg pre- emergence, isoproturon at 2.5 kg at two leaf stage clodinafop-propargyl at 0.06 kg at one tiller stage
11	Tri-allate at 2.24 kg plus trifluralin at 0.96 kg pre- emergence, isoproturon at 1.5 kg plus simazine at 0.25 kg at two leaf stage
12	Fenoxaprop-P-ethyl at 0.069 kg at two leaf stage
13	Fenoxaprop-P-ethyl at 0.069 kg at one tiller stage
14	Clodinafop-propargyl at 0.06 kg at two leaf stage
15	Clodinafop-propargyl at 0.06 kg at one tiller stage

Experimental diary:

20-Sep-95 : B : Ploughed and furrow pressed.
22-Sep-95 : B : Rolled.
10-Oct-95 : B : Gramoxone 100 at 3.0 l in 260 l.
12-Oct-95 : B : Rotary harrowed.

96/R/WW/6

Experimental diary:

13-Oct-95 : B : Rotary harrowed Mercia, dressed Sibutol, drilled at 380 seeds per m².
17-Oct-95 : B : Draza at 5.5 kg.
18-Oct-95 : T : HERBCIDE 5, 7, 8, 9, 10, 11: Portman Trifluralin at 2.0 l in 220 l.
 : T : HERBCIDE 6: Stomp 400 SC at 5.0 l in 220 l.
19-Oct-95 : T : HERBCIDE 4, 7, 8, 9, 10, 11: Avadex BW Granular at 22.4 kg.
10-Jan-96 : T : HERBCIDE 11: Atlas Simazine at 0.5 l with Stefes IPU at 3.0 l in 220 l.
 : T : HERBCIDE 12: Cheetah Super at 1.25 l in 220 l.
 : T : HERBCIDE 3, 4, 5, 6, 7, 8, 9, 10: Stefes IPU at 5.0 l in 220 l.
 : T : HERBCIDE 14: Topik at 0.25 l in 220 l.
02-Apr-96 : T : HERBCIDE 8: Stefes IPU at 4.2 l in 220 l.
 : T : HERBCIDE 9, 13: Cheetah Super at 1.25 l in 220 l.
 : T : HERBCIDE 10, 15: Topik at 0.25 l in 220 l.
18-Apr-96 : B : 34.5% N at 586 kg.
12-Jun-96 : B : Monicle at 1.0 l in 300 l.
20-Aug-96 : B : Combine harvested.

Previous crops: W. wheat 1994 and 1995.

NOTE: Black-grass and wheat plants were counted in December. Black-grass heads were counted in June.

96/R/WW/6

GRAIN TONNES/HECTARE

***** Tables of means *****

HERBCIDE

1	5.17
3	7.03
4	7.48
5	7.54
6	7.33
7	7.19
8	7.19
9	7.01
10	7.25
11	6.99
12	6.13
13	5.78
14	6.99
15	6.15

Mean 6.69

*** Standard errors of differences of means ***

HERBCIDE

0.256 min.rep
0.222 max-min

HERBCIDE

max-min 1 v any of the remainder
min.rep Any of the remainder

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	43	0.362	5.4
GRAIN MEAN DM%	88.2		
PLOT AREA HARVESTED	0.00219		